


Northwest Fishery Resource Bulletin



Coho Salmon Escapement to the Skagit River Estimated Using a Mark-Recapture Method: 1988

By

Robert H. Conrad

Northwest Indian Fisheries Commission

Robert A. Hayman

Skagit System Cooperative

Eric A. Beamer

Skagit System Cooperative,

and

Pamela J. Goddard

Northwest Indian Fisheries Commission

Project Report Series No. 8

Northwest Fishery Resource Bulletin

Project Report Series

The Northwest Fishery Resource Bulletin presents the results of investigations carried out by the Washington Dept. of Fish and Wildlife, Western Washington Treaty Tribes, and/or the Northwest Indian Fisheries Commission that are deemed of sufficient interest to be made available to the scientific community and the public.

The Project Report Series is designed to report on the results of research and data collection projects or significant work in progress that may have immediate useful applications.

The contents of this report may be reprinted with the permission of the authors. Reference to the source is requested.

This report series sponsored by the:


Washington Department of Fish and Wildlife

Western Washington Treaty Tribes

Northwest Indian Fisheries Commission

Inquiries should be addressed to:

Northwest Indian Fisheries Commission
6730 Martin Way East
Olympia, WA 98516-5540
Phone: (360)-438-1180



Printed on recycled paper

Coho Salmon Escapement to the Skagit River Estimated Using a Mark-Recapture Method: 1988

by

Robert H. Conrad: Northwest Indian Fisheries Commission¹,
Robert A. Hayman: Skagit System Cooperative²,
Eric A. Beamer: Skagit System Cooperative,
and
Pamela J. Goddard: Northwest Indian Fisheries Commission

Northwest Fishery Resource Bulletin
Project Report Series No. 8

May 1998

¹ Northwest Indian Fisheries Commission, 6730 Martin Way E., Olympia WA 98516

² Skagit System Cooperative, PO Box 368, LaConner WA 98257

This page left blank intentionally.

ABSTRACT

Since 1965, an index live-count method has been used to annually estimate the number of coho salmon in the escapement to the Skagit River. The accuracy and precision of the estimates from this method have never been critically examined. A five-year project to examine alternative methods of estimating the number of wild coho salmon in the escapement to the Skagit River was begun in 1986. In addition to the index live-count method, three other methods of estimating the coho salmon escapement to the Skagit River were examined: (1) a mark-recapture method; (2) a redd-count expansion method; and (3) a method based on estimates of the proportional contribution of hatchery-produced coho salmon to the total escapement. **This report documents the results of the mark-recapture portion of the project for 1988.**

In 1988, coho salmon were captured with a beach seine at river mile 35 near the town of Lyman on the Skagit River from 12 September through 15 November. A total of 1,970 coho salmon were tagged with a jaw tag and marked with opercula punches. Tags were recovered during surveys designed to sample randomly the coho salmon escapement. Tag recovery samples were collected at 13 areas in the Skagit River drainage: Marblemount Hatchery; Baker River trap; spawning grounds in the Middle Skagit, Upper Skagit, Lower Sauk, Middle Sauk, Upper Sauk, Suiattle, Cascade, Nookachamps, and Carpenter sub-basins; and in commercial or test fisheries. A total of 24,552 coho salmon were examined of which 24,028 fish were considered in-sample and 524 were not considered part of the population subject to tagging.

A total of 464 tagged or marked coho salmon were recovered during in-sample surveys. The tag recovery data indicate that approximately 2% of the coho salmon migrating through the lower Skagit River tagging area were caught and tagged. About 3% of the coho salmon returning to Marblemount Hatchery and the Baker River trap had jaw tags compared to about 2% of the fish in the samples from most of the sub-basin spawning grounds above the tagging area, but this difference was not significant. The tag recovery data indicate that some coho salmon from spawning areas substantially downstream of the tagging site were present in the tagging area. There was one tag recovered in 1,257 coho salmon examined (0.08%) during spawning ground surveys in the Nookachamps and Carpenter sub-basins.

The estimated abundance of coho salmon in 1988 was 70,751 fish with a 95% confidence interval of 64,614 to 78,516 fish. This estimate is for the number of coho salmon migrating through the tagging area after tagging began on 12 September. It includes all coho salmon bound for spawning areas above the tagging area and an unknown fraction of the salmon from spawning areas in the Nookachamps and Carpenter sub-basins. This abundance estimate was very precise (CV = 4.4%) because of the large number of fish examined for tags during in-sample surveys. To restrict the estimate to spawning areas in the Middle Skagit sub-basin and spawning areas above it, adjustments were made to the number of tags released. Using the adjusted number of tags released, the estimated abundance for this more restricted area was 70,392 coho salmon. **The total return of coho salmon to Skagit Bay in 1988 is estimated to be 86,265 fish. There were an estimated 61,283 naturally-spawning coho salmon in the escapement to Skagit River spawning grounds: 58,029 fish were estimated to have reached upstream spawning grounds and 3,254 coho salmon were estimated for lower river (Nookachamps and Carpenter sub-basins) spawning grounds (see summary table on the next page).**

Table summarizing the total return of coho salmon to Skagit Bay in 1988 by its major components.

Component	Number of Fish
Total Terminal Run Size	86,265
Marblemount Hatchery	8,783
Baker River Hatchery	1,500
Commercial Fishery Catches	12,079
<u>Test Fishery Catches</u>	<u>2,620</u>
Subtotal	24,982
Wild Escapement	
Upstream Areas	58,029
<u>Lower Areas</u>	<u>3,254</u>
Subtotal	61,283
Sport Catch ^a	1,050

^a An unknown portion of the sport catch should be subtracted from the wild escapement and the remainder added to the total terminal run size.

TABLE OF CONTENTS

	<u>PAGE</u>
ABSTRACT	i
LIST OF TABLES	v
LIST OF FIGURES	viii
INTRODUCTION	1
METHODS	3
Tagging Methods	3
Beach Seining	3
Tagging Procedures	5
Tag Recovery Surveys	5
Marblemount Hatchery	5
Baker River Trap	6
In-River Commercial and Test Fisheries	8
Spawning Grounds	8
Tributary Traps	10
Abundance Estimates	10
Petersen Estimation Model	11
Darroch's Stratified Estimation Model	11
Definition of Strata	11
Testing ρ and π	12
Selection of Estimation Models	12
Allocating Marked-Only Fish to Releases Strata	13
Additional Analyses	13
Migratory Timing to Major Recovery Areas	13
Analyses of Sex and Length Composition Data	14
RESULTS	16
Tag Releases	16
Tag Recoveries	21
Marblemount Hatchery	21
Baker River Trap	21
Commercial and Test Fishery Samples	24
Middle Skagit Sub-basin	24
Upper Skagit Sub-basin	24
Lower Sauk Sub-basin	25
Middle Sauk Sub-basin	25
Upper Sauk Sub-basin	25
Suiattle Sub-basin	25

TABLE OF CONTENTS (continued)

	<u>PAGE</u>
Other Spawning Ground Surveys	25
Out-of-System Recoveries	26
Abundance Estimates	26
Additional Analyses	28
Timing of Migrations to Major Recovery Areas	28
Length and Sex Composition Analyses	28
Conclusions	33
DISCUSSION	36
Population was Closed	36
Area Encompassed by the Estimates	36
Estimate of the Number of Tagged Fish “Lost” to Areas Downstream of the Tagging Area	37
All Coho Salmon Have an Equal Probability of Capture During Tagging or the Recovery Sample is a Simple Random Sample of the Population	38
Tagging Does Not Affect the Catchability of an Animal	38
Animals Do Not Lose Their Tags Between the First and Second Samples	39
All Tagged Animals are Reported in the Second Sample	39
There are No Mortalities Due to Tagging	39
CONCLUSIONS	40
ACKNOWLEDGMENTS	42
REFERENCES CITED	43
APPENDIX A - Summary tables of sample data for 1988	45
APPENDIX B - Details of abundance estimates generated for 1988	66

LIST OF TABLES

<u>TABLE</u>	<u>PAGE</u>
1. Number of coho salmon tagged each day and number of in-sample tag recoveries from each day's release for the Skagit River, 1988	17
2. Summary of the number of tag recoveries (#) from each release stratum in each major recovery area and the results of testing recovery percentages (π) for equality among release strata, 1988	19
3. Summary of the percentage of tagged or marked coho salmon found in each recovery area during in-sample surveys of the Skagit River, 1988	22
4. Average day of release (DOR) and average number of days between release and recovery (DBET) for coho salmon tagged and recovered in the Skagit River, 1988	23
5. Summary of estimates of the number of coho salmon in the Skagit River escapement using data from each major recovery area, 1988	27
6. Summary of the number of coho salmon returning to Skagit Bay in 1988	41

<u>APPENDIX TABLE</u>	<u>PAGE</u>
A-1. Summary of coho salmon escapement samples collected at Marblemount Hatchery in 1988	46
A-2. Summary of coho salmon escapement samples collected at Baker River trap in 1988	47
A-3. Summary of coho salmon catch samples collected from the commercial fishery in the Skagit River, 1988	48
A-4. Summary of coho salmon catch samples collected during test fisheries in the Skagit River, 1988	49
A-5. Summary of coho salmon escapement samples from the Middle Skagit sub-basin collected during spawning ground surveys by Skagit System Cooperative crews and at a trap on Careys Creek, 1988	50

LIST OF TABLES (continued)

<u>APPENDIX TABLE</u>	<u>PAGE</u>
A-6. Summary of coho salmon escapement samples from the Middle Skagit sub-basin collected at a trap on Etach Slough operated by Skagit System Cooperative, 1988	51
A-7. Summary of coho salmon escapement samples from the Upper Skagit sub-basin collected during spawning ground surveys by Skagit System Cooperative crews, 1988	53
A-8. Summary of coho salmon escapement samples from the Lower Sauk sub-basin collected during spawning ground surveys by Skagit System Cooperative crews, 1988	54
A-9. Summary of coho salmon escapement samples from the Middle Sauk sub-basin collected during spawning ground surveys by Skagit System Cooperative crews, 1988	55
A-10. Summary of coho salmon escapement samples from the Upper Sauk sub-basin collected during spawning ground surveys by Skagit System Cooperative crews, 1988	56
A-11. Summary of coho salmon escapement samples from the Suiattle sub-basin collected during spawning ground surveys by Skagit System Cooperative crews and at a trap on Suiattle Slough, 1988	57
A-12. Summary of coho salmon escapement samples from the Nookachamps sub-basin collected during spawning ground surveys by Skagit System Cooperative crews and at traps on the East Fork and West Fork of Nookachamps Creek, 1988	58
A-13. Summary of coho salmon escapement samples from the Carpenter sub-basin collected during spawning ground surveys by Skagit System Cooperative crews and at a trap on Carpenter Creek Slough, 1988	60
A-14. Summary of coho salmon escapement samples from the Cascade sub-basin collected during spawning ground surveys by Skagit System Cooperative crews, 1988	61
A-15. CPUE (catch per beach seine set) of coho salmon bound for major recovery areas in the Skagit River, 1988	62

LIST OF TABLES (continued)

<u>APPENDIX TABLE</u>	<u>PAGE</u>
A-16. Summary of the number of tag releases and number of in-sample tag recoveries by length for male and female coho salmon tagged in the lower Skagit River, 1988	63
A-17. Daily summary of the numbers of coho salmon tagged in the lower Skagit River and recovered during in-sample surveys, by sex, release condition, and maturity classification, 1988	64
A-18. Summary of the estimated number of tags from areas downstream of the tagging area in the lower Skagit River, 1988	65

LIST OF FIGURES

<u>FIGURE</u>	<u>PAGE</u>
1. Map of the Skagit River system showing the location of the areas where tag recovery surveys were conducted	2
2. Map of the area of the Skagit River where coho salmon were captured and tagged, 1986-1990	4
3. Schematic of the sampling procedure used to process coho salmon for tag examination at Marblemount Hatchery	7
4. Commercial fishery areas of the Skagit River as designated by WDFW	9
5. Percent of tags recovered during in-sample surveys from each day of release for coho salmon tagged in the Skagit River, 1988	18
6. Catch-per-unit effort of coho salmon by the beach seine in the lower Skagit River tagging area by day and for each ten-day period, 1988	29
7. Beach seine catch-per-unit effort (CPUE) of coho salmon bound for major Skagit River tag recovery areas in 1988	30
8. Comparison of length frequencies of male and female coho salmon tagged in the lower Skagit River, 1988	31
9. Comparison of length frequencies of coho salmon that were tagged but not recovered to those that were tagged and recovered, for males and females, 1988	32
10. Sex and length composition, by release period, of coho salmon tagged in the lower Skagit River, 1988	34

INTRODUCTION

The Skagit River is the largest river system in the Puget Sound region. It has 162 miles of mainstem river and its headwaters are in Canada (Figure 1). This system is one of the largest producers of coho salmon (*Oncorhynchus kisutch*) in northern Puget Sound. Coho salmon from the Skagit River are caught in fisheries from Northern California to Southeast Alaska and are a major contributor to fisheries in the inside marine waters of Georgia Strait and Puget Sound (PFMC 1992). The Skagit River is managed for natural production of coho salmon (subsequently referred to as wild coho salmon). In years when the numbers of wild coho salmon projected to return to the Skagit River are small, fisheries from Cape Falcon, Oregon to the US/Canada border have been constrained to protect these fish (PFMC 1986, pg. III-9; and PFMC 1988, pg. III-11). Accurate annual assessments of stock status are required for coho salmon from the Skagit River because this stock can affect the management of fisheries over such a large geographic area. This ensures that fisheries are not unnecessarily restricted during years when there is not a conservation problem and prevents over-harvest of wild coho salmon from the Skagit River during years of small returns. An important component of the information needed to accurately assess the status of wild coho salmon from the Skagit River is an annual estimate of the number of coho salmon in the spawning escapement. Spawning escapement, as used in this report, refers to the number of adult coho salmon which are present in all natural spawning areas of the Skagit River and have the potential to spawn in these areas. It does not include coho salmon returning to Marblemount Hatchery or to the release site for hatchery-produced coho salmon at the Baker River dam.

Since 1965, the Washington Department of Fish and Wildlife (WDFW) has used an index live-count method to annually estimate the escapement of coho salmon to the Skagit River (Flint 1983). The accuracy and precision of the estimates from this method have not been critically examined. A five-year project to examine alternative methods of estimating the number of wild coho salmon in the spawning escapement to the Skagit River was begun in 1986. This project was conducted by the Skagit System Cooperative (SSC) in cooperation with personnel from WDFW and Puget Power and Light. Three methods of estimating the spawning escapement of coho salmon to the Skagit River were examined: (1) a mark-recapture method; (2) a redd-count method; and (3) a method based on estimates of the proportional contribution of hatchery-produced coho salmon to the total escapement.

This report is the third in a series of reports that will document the studies conducted from 1986 through 1990 which examined different methods for estimating the escapement of coho salmon to the Skagit River. The 1986 and 1987 studies are summarized, respectively, in Conrad et al. (1997 and 1998). **This report summarizes the data and documents the results of the mark-recapture portion of the project for 1988.** Reports documenting the results for the other years that tagging was conducted (1989 and 1990) and the other methods of estimation will follow. Some summary data from the other years of the study are used to support some of the assumptions required for the analysis of the tagging data from 1988. These data are documented in Conrad et al. (1997).




Figure 1. Map of the Skagit River system showing the location of the areas where tag recovery surveys were conducted.

METHODS

The description of methods is divided into four sections. The first section describes the methods used to capture coho salmon for tagging and the tagging procedure. The second section describes the surveys used to recover tags. This includes a description of the survey procedures for each of the tag recovery areas. Section three summarizes the statistical procedures used to estimate the abundance of coho salmon from the tag release-and-recovery data. The last section describes some miscellaneous analyses conducted to examine migration timing and the sex and length composition of the coho salmon that were sampled.

Tagging Methods

Beach Seining:

Coho salmon were captured for tagging using a beach seine operated by a five-man crew. Seining was conducted at a single site at about river mile 35 near the town of Lyman on the Skagit River (Figure 2). A beach seine that was 456' long by 20' deep was used to capture coho salmon. The seine had two wings: one was 90' long and made of 3.5" knotless nylon and the other was 330' long and made of 2.75" monofilament. The net had a 36' bunt made of 2" knotless seine material. Cork spacing was 8" on the bunt and two feet on the rest of the net. The leadline was hung with 15 lb per 60' of net. Modifications in net dimensions occurred whenever the seine was damaged. Due to heavy use, the leadline was rehung about every four fishing days and the monofilament was replaced after every eight to ten fishing days.

A boat was used to set the beach seine. One end of the seine was held by two crew members on a gravel bar while the boat backed away from the shore and the net was set off the bow of the boat. When the entire net was out, the boat-end of the net was towed downstream. The other end of the net was attached to a four-wheel drive truck and driven slowly downstream. Care was taken to prevent the shore-end of the net from getting ahead of the boat because fish tended to lead away from the shore and around the boat. During the drift, a seine plunger (a long pole with a cup on the end) was slammed into the water periodically to drive fish away from the river-end of the net and toward the shore. At a pre-designated point, the boat returned to the gravel bar. Upon reaching the shore, the boat-end of the net was attached to the back of a second four-wheel drive truck. Both trucks then pulled the net up the gravel bar, perpendicular to the river, until only the bunt end of the net was in the water. The five-man crew then pulled the bunt in by hand until the leadline was on shore while the cork line and ends were cradled by the crew.




Figure 2. Map of the area of the Skagit River where coho salmon were captured and tagged, 1986-1990.

Tagging Procedures:

Coho salmon were removed from the bunt and placed into either of two net pens adjacent to the capture site. All other species were counted and returned to the river. The pens were 3' by 5' by 5', constructed of PVC, and covered with 0.5" knotless nylon mesh. Each coho salmon was taken from a net pen by a sampler wearing cotton gloves and placed on a V-shaped measuring board lined with high-density foam. A sequentially-numbered hog ring was clamped around the lower left mandible of each fish using a pair of hog-ring pliers and a 3/8" hole was punched in the rear center of each gill operculum with a paper hole-puncher. The fork length (measured to the nearest cm), sex of the fish, any external marks, and a qualitative assessment of maturity (bright, blush, or dark) were recorded for each fish with the date and tag number. Each tagged salmon was held gently in the water until its equilibrium was regained before being released. If a tagged fish did not swim away or appeared to be injured it was given a condition rating of "X-". Fish that swam away normally were given a condition rating of "X". If a fish was especially vigorous when released a condition rating of "X+" was assigned. Fish with severe physical impairments (e.g., 50% scale loss, torn opercula, deep predator wounds) were released untagged. These included jack coho salmon (male salmon under 30 cm in length) which generally gilled in the net and were unfit for tagging.

Tag Recovery Surveys

Only tags recovered during surveys designed to randomly sample the coho salmon escapement were used for the abundance estimates. These are referred to as **in-sample recoveries**. Tag recovery surveys were conducted by sampling: (1) all fish spawned, surplused, or otherwise sacrificed at Marblemount Hatchery; (2) all fish caught at the fish trap at Baker River dam; (3) the catch by the in-river commercial fishery; (4) all test fishery catches; (5) every reachable and identifiable dead coho salmon found during spawning grounds surveys; and (6) every coho salmon caught in traps operated on: Fisher Creek and Carpenter Creek Slough (tributaries to Carpenter Creek); the East Fork of Nookachamps Creek; Hansen Creek (a tributary to the Middle Skagit sub-basin); and Barnaby Slough (a tributary to the Upper Skagit sub-basin). During each survey or day of trap operation, the date, number of coho salmon inspected for tags, number of tagged or marked (with the opercula punches) fish found, and tag numbers of all coho salmon recovered with legible jaw tags were recorded.

Marblemount Hatchery:

Samples were collected by three different methods at WDFW's Marblemount Hatchery: spawned fish, surplused fish, and pond mortalities. After any processing, hatchery personnel sorted the fish from these groups into separate bins for tagged/marked and unmarked fish. SSC crews then re-checked these bins for coho salmon with tags or marks. The date of sampling, number of coho salmon inspected for tags, number of tagged or marked fish found, and tag numbers of all coho salmon recovered with legible jaw tags were recorded.

Coho salmon were spawned at Marblemount Hatchery to meet specific egg-take goals. Spawning was conducted when the portion of the run from which eggs were desired was present and there were large numbers of fish in the holding ponds. Hatchery personnel selected fish for spawning and sorted them into the bins after spawning for SSC crews to examine. Surplussed fish were those in excess of the spawners needed for eggs. Surplus coho salmon were periodically sacrificed and sorted into the bins. The holding pond was periodically surveyed for mortalities and any dead coho salmon were removed and sorted into the bins. A schematic of the Marblemount Hatchery sampling procedure is shown in Figure 3.

Except for the pond mortalities, hatchery personnel selected the coho salmon for the other two groups, spawned and surplussed, according to a visual assessment of the fish and the timing of the return to Marblemount Hatchery. Therefore, these fish were not strictly sampled at random and the percentage of tagged fish in these samples might have been influenced by the selection process. However, since all coho salmon returning to the hatchery were sampled, the Marblemount Hatchery sample was a census and the sample total for the entire spawning season provided the best estimate of the percentage of tagged coho salmon at Marblemount Hatchery.

Baker River Trap:

A fish trap at Baker River dam caught all upstream migrating salmon. All coho salmon caught at the trap were examined. Fish caught in the trap were crowded into a brail and several removed at a time onto a sorting table. Each coho salmon was examined for a tag or mark. The sample date, condition, and tag number (when legible) were recorded for any jaw-tagged or opercula-punched coho salmon. After all live fish in the brail were removed, the racks and screen of the trap were searched for dead fish. Therefore, identically to the Marblemount Hatchery sample, the Baker River trap sample was a census and the sample total for the entire spawning season provided the best estimate of the percentage of tagged coho salmon at the Baker River trap.

The Baker River stock is one of the earliest returning coho salmon stocks to the Skagit River. Coho salmon were counted at the Baker River trap before tagging began in the lower river during two years of the study. In the other years of the study, coho salmon were counted at the trap so soon after tagging was initiated that we assumed some fish had migrated past the tagging site before tagging had begun and, therefore, were not subject to capture. Since these early-arriving fish were not subject to tagging, we excluded them from the number of fish examined for tags that was used for the population estimates (i.e., they were not considered in-sample). We examined the number of days between release and recapture for all coho salmon recovered at the Baker River trap during the five years of tagging. The minimum travel time (number of days between being tagged and released in the lower river and recovered at Baker River trap) observed during the study years was four days (Conrad et al. 1997). Therefore, all fish counted at the Baker River trap prior to four days after tagging had begun were excluded from the in-sample survey.




Figure 3. Schematic of the sampling procedure used to process coho salmon for tag examination at Marblemount Hatchery.

In-River Commercial and Test Fisheries:

Tag recovery samples from the commercial catch were collected in conjunction with routine commercial catch sampling activities. The Skagit River is divided into statistical areas for commercial catch regulation (Figure 4). To allow tag recovery samples from the catch to be analyzed by area of capture, all major salmon buyers were instructed to place catches from each statistical area into separate bins. This occurred during the fishery for coho salmon in the Upper Skagit River; there was no commercial fishery for chum salmon in the Upper Skagit River in 1988. Most samples were allocated to sub-areas (78D-2, 78D-3, etc.) within Area 78D except for incidental catches of coho salmon during the steelhead fishery in the Upper Skagit River which were not sorted by area. The distribution of these samples among the sub-areas was used to allocate the entire

A test fishery was conducted each year by an SSC crew to provide an in-season assessment of the size of the coho salmon run. In 1988, test fisheries were conducted in: Area 2; Spudhouse; Blakes; and Bay and Jetty in Skagit Bay (Figure 4). Drift and set gill nets used at the test fish sites had mesh sizes ranging from 5" to 6". Hayman (1996) describes the test fishing procedures in detail. All coho salmon caught during the test fishery were inspected for tags or marks.

Both WDFW and tribal commercial catch and hatchery samplers in areas outside of the Skagit River were notified to look for jaw tags from the Skagit River study. These recoveries allowed us to assess the degree of out-of-system straying for coho salmon tagged in the mainstem of the Skagit River.

Spawning Grounds:

Tag recovery surveys of the spawning grounds were conducted in conjunction with surveys to estimate the coho salmon escapement using redd counts (Conrad et al. 1993). For the redd-count method, the Skagit River system was stratified into the nine sub-basins listed by Johnson (1986): Carpenter; Nookachamps; Middle Skagit; Upper Skagit; Lower Sauk; Middle Sauk; Upper Sauk; Suiattle; and Cascade (Figure 1). Stream sections in each sub-basin were surveyed from one to 18 times during the spawning period for coho salmon. In 1988, about 25% of the total length of potential spawning habitat in the Skagit River was surveyed (Conrad et al. 1993). During spawning ground surveys, any coho salmon carcasses observed were sampled for jaw tags and opercula marks. Gill opercula of untagged carcasses were carefully inspected for marks or healed marks. A healed (regenerated) mark was evident as a perfectly round discoloration on the gill cover that was lighter in color than the surrounding opercular tissue. Occasionally a carcass could not be sampled because of a missing head due to advanced decomposition or removal by predators. Unsourced carcasses were tallied during each survey. The date, survey location, number of coho salmon carcasses sampled, number of tagged or marked fish recovered, and tag numbers of all coho salmon recovered with legible jaw tags were recorded during these surveys. The caudal fin of all sampled carcasses was removed to prevent the carcass from being sampled again during subsequent surveys.




Figure 4. Commercial fishery areas of the Skagit River as designated by WDFW.

Tributary Traps:

There were six tributary traps operated in the Skagit system in 1988: four were operated by SSC and two were operated by WDFW. SSC operated traps on: Carpenter Creek Slough (a tributary to the Carpenter sub-basin); Etach (or Red Cabin) Slough (a tributary to the Middle Skagit sub-basin); the East Fork of Nookachamps Creek; and the West Fork of Nookachamps Creek (tributaries to the Nookachamps sub-basin). The two traps operated by WDFW were part of a habitat enhancement study and were located on Careys Creek (a tributary to the Middle Skagit sub-basin) and Suiattle Slough (a tributary to the Suiattle sub-basin). The traps on Carpenter Slough and Etach Slough were wooden weirs that blocked the entire creek and funneled fish into a live box. The traps on the East and West Forks of Nookachamps Creek were wire-mesh hoop traps that blocked only a portion of the creek but had visual lead nets attached that stretched from bank to bank. All traps were located in areas that had easy accessibility, a section of relatively straight stream channel with a low gradient, and a stable substrate.

All traps were checked and cleaned at least twice daily. A knotless-nylon dip net was used to move the trapped coho salmon into a 30-gallon plastic container filled with water. All coho salmon caught were examined for tags or marks and then released upstream. A Petersen disk tag and a unique operculum punch (i.e., a punch pattern different from that used in the main-river tagging) were placed on all coho salmon released above the traps. The trap crews also recovered tags at the weirs from spawned-out carcasses which had washed downstream from the spawning areas (called rack recoveries). The caudal fin was cut off all rack recoveries. The date, number of coho salmon sampled, number of tagged or marked fish recovered, and tag numbers of all coho salmon recovered with legible jaw tags were recorded.

Abundance Estimates

Two different mark-recapture models were used to estimate the number of coho salmon passing through the tagging area in the lower Skagit River, the Petersen estimation model and Darroch's stratified estimation model. When tagging and recovery occur over an extended time period, such as occurred in this study, it is not uncommon to observe temporal changes in: (1) the probability of capture of fish in the target population; and/or (2) the probability of finding a tagged fish during tag recovery surveys. When such changes occur the Petersen model is often not the appropriate estimation model. Seber (1982) describes a series of χ^2 tests to determine whether the data are consistent with a Petersen estimate. Specifically, the tests determine whether the data are consistent with the following four conditions: (1) there was uniform recovery of tags across the tag recovery strata; (2) there was uniform tagging across the tag release strata; (3) there was complete mixing of the population between tagging and recovery; and (4) the expected number of tags recovered in each stratum was proportional to the number of unmarked individuals present.

Eames et al. (1981, 1983) describe the exact form of these tests for a study similar to ours in both the study design and estimation procedures. They captured chum and coho salmon in marine areas immediately in front of the mouths of major river systems in Puget Sound and

tagged the fish with jaw tags. Tags were recovered during surveys of spawning grounds throughout these river systems. We followed procedures similar to those described by Eames et al. (1981, 1983) to determine the appropriate estimation model.

Petersen Estimation Model:

The simplest and most commonly used model for estimating abundance from mark-recapture data is the Petersen model. Conrad et al. (1997) discuss the necessary assumptions for the Petersen model as implemented for this study.

Robson and Regier (1964) recommend that a Petersen estimate include a minimum of seven tag recaptures to ensure that the bias of the estimate is negligible. Therefore, we estimated abundance from the tagging data only when there were at least seven recaptures of tagged or marked coho salmon from a recovery area. Chapman's unbiased form of the Petersen estimate (Seber 1982) was used to estimate abundance. Conrad et al. (1997) describe the model and the procedures used to estimate 95% confidence intervals. For any Petersen-type estimator (including Darroch's stratified estimator), the abundance estimate depends upon ρ , the proportion of the population tagged. The proportion of tags in the second (recovery) sample provides an estimate of ρ . Generally, as $\hat{\rho}$ becomes smaller the estimated abundance becomes larger for a given number of tags released.

Darroch's Stratified Estimation Model:

Darroch (1961) developed a stratified population model for open populations that is not predicated on constant probabilities of capture or recovery. The necessary assumptions for this model are discussed in Seber (1982) and summarized by Conrad et al. (1997). Conrad et al. (1997) also describe the model and its application to the tag release-and-recovery data collected for this study.

Definition of Strata:

Two different tag recovery percentages were examined to help define tag release and tag recovery strata. To determine if the probability of finding a tagged fish in recovery samples was different among recovery locations or among different time periods at the same location, the percentages of tags in recovery samples (ρ as defined previously) were compared. The percentages of tags recovered from releases during specific time strata, π , were compared to determine if there were differences in the probability of recovering fish tagged during different segments of the release period. For these tests it was necessary to define temporal strata for both the tag release data and the tag recovery data from each recovery area.

Tag release strata were established by dividing the release data into four to six strata with about an equal number of days of tagging in each stratum. The percentages of tagged fish recovered from each release stratum (π) were tested to determine if they were equal. If a significant difference was found ($P \leq 0.10$) additional χ^2 tests were conducted to more

precisely define the release strata by pooling adjacent strata which did not have significantly different π .

Three different criteria were used to establish tag recovery strata: (1) number of days of sampling; (2) number of tags recovered; and (3) number of fish examined for tags. Initially, two recovery strata were defined by dividing the data so there were approximately equal numbers of the criteria (days surveyed, number of tags, or number of fish examined) in each stratum. The percentages of tagged fish in each recovery stratum (ρ) were tested to determine if they were equal among recovery strata for each stratification criteria. If a significant difference was found ($P \leq 0.10$) additional χ^2 tests were conducted within the initial strata to more precisely define the recovery strata.

Testing ρ and π :

Tests were conducted to determine if there were significant differences in tag recovery percentages (either ρ or π) between different samples or groups of fish (e.g., between surveys conducted by SSC and WDFW, or between samples collected during different time periods, or between samples collected at different locations, or between male and female coho salmon). When the expected number of tag recoveries for each group in a comparison was five or greater, a standard χ^2 test (Conover 1980) was used to test for differences in tag recovery percentages (ρ or π). If the number of tag recoveries was insufficient for a χ^2 test (one or more cells with expected frequencies less than five) and there were only two release strata or recovery locations to compare, Fisher's exact test (Conover 1980) was used. Otherwise, an approximate randomization test (ART) was conducted (Noreen 1989). An approximate randomization test is a computer-intensive method of testing whether the data in a contingency table are similar. It is similar to Fisher's exact test but uses a computer to repeatedly resample the data and approximately estimate the probability of observing the configuration of the data in the table (under the null hypothesis that the samples are from the same population).

Selection of Estimation Models:

If we assume that coho salmon bound for each recovery area are randomly sampled as they migrate through the lower river tagging area, the recovery data (number of tagged or marked fish found and number of fish examined) from each recovery area can be used to estimate ρ , the percentage of the population that was tagged. If the hypothesis of equal $\hat{\rho}$ among recovery areas was not rejected ($P > 0.10$), the tag recovery data from the different areas were pooled. The pooled data were then used in the tests to determine if the tag release-and-recovery data were consistent with the Petersen model. We feel that the variation in $\hat{\rho}$ among the recovery areas generally reflects sampling variation in the recovery areas. The number of carcasses examined for tags was relatively small for some recovery areas. In some cases, all samples were collected from a relatively discrete area within the general recovery area which could influence the proportion of tagged carcasses present. Generally, the areas with greatly different recovery percentages (more than a 0.5% difference from the major recovery areas) had less than seven tag recoveries each. The different population estimates that were

generated using the data from different recovery areas (or pooled recovery areas) were usually not significantly different from each other. Therefore, we selected the estimate with the smallest coefficient of variation as the “best” estimate of abundance for each year.

The model used to estimate abundance, simple Petersen or Darroch’s stratified, was determined by the results of the tests for the consistency of the data. The four χ^2 tests used to determine consistency are described by Seber (1982) and by Eames et al. (1981, 1983).

Allocating Marked-Only Fish to Release Strata:

From 12% to 24% of the in-sample recoveries each year had a tag with an illegible number or had no tag and were identified as tagged fish by the opercula punches. The release stratum for these fish was unknown and had to be estimated for the stratified estimator. Marked fish with missing or illegible tags were allocated to release strata within a recovery area based on the proportional distribution of legible tags from each release stratum (Conrad et al. 1997). This assumes that tag loss or tag illegibility is a random process and that coho salmon tagged during each release stratum have equal rates of tag loss, therefore, fish with missing or illegible tags are assumed to have a similar distribution for stratum of release as fish with legible tags. If tag loss (or a tag becoming illegible) is a time dependent process, then fish tagged during the earlier release strata might be expected to have higher rates of tag loss and this assumption would not be true. Eames et al. (1981, 1983) used procedures similar to ours to allocate fish recovered with missing tags to release strata in their study. Errors in the assignment of marked-only fish to release strata affect only the Darroch estimate.

Additional Analyses

Several additional analyses of the data collected during tagging and tag recovery surveys were conducted. These included analyses to determine the timing of the migration of different spawning groups through the tagging area and analyses of sex and length composition data. These analyses were not required for the abundance estimates but were conducted to describe the characteristics of the annual return of coho salmon to the Skagit River during the study years.

Migratory Timing to Major Recovery Areas:

The timing of coho salmon migrating through the lower river tagging area was estimated from an analysis of the release dates of the tags recovered in each major recovery area (excluding commercial and test fisheries). Only areas with ten or more legible tag recoveries were included in the analyses. Ten, 10-day time periods were defined for the migratory timing calculations: (1) 1 September to 10 September; (2) 11 September to 20 September; (3) 21 September to 30 September; (4) 1 October to 10 October; (5) 11 October to 20 October; (6) 21 October to 30 October; (7) 31 October to 9 November; (8) 10 November to 19 November; (9) 20 November to 29 November; and (10) 30 November to 9 December.

Catch-per-unit effort (CPUE) by the beach seine used to capture coho salmon for tagging was used to describe the timing of the run through the tagging area in the lower river. CPUE was calculated for each 10-day period as the total number of coho salmon caught divided by the total number of beach seine sets (catch per set). The number of tags recovered in each major recovery area from each of the release periods was used to estimate the CPUE of coho salmon bound for these areas. The CPUE of coho salmon from recovery area j during release period i was estimated by:

$$\hat{\omega}_{ij} = \frac{r_{ij}}{f_i} \quad [6]$$

where $\hat{\omega}_{ij}$ = the estimated CPUE of coho salmon from recovery area j during release period i ,
 r_{ij} = the number of tags recovered in area j that were released during period i , and
 f_i = the number of beach seine sets made during period i .

For each area analyzed, the CPUE estimated for each 10-day period was summed across all ten time periods to estimate a season total CPUE of coho salmon bound for that recovery area. The estimated CPUE of coho salmon from recovery area j during time period i was converted to the percentage of this season total CPUE for recovery area j to describe migratory timing (Mundy 1982). These data were then graphed so that the migratory timing patterns for the major recovery areas could be compared.

Analyses of Sex and Length Composition Data:

Significant differences in the probability of recovering coho salmon tagged during different release periods (π) were found at some recovery locations in 1988. Temporal trends in the probability of recovery could be due to changing environmental conditions at the tagging site which influenced the probability of capture. For example, high and low water conditions may have influenced the effectiveness of the beach seine used to capture fish in the tagging area. Under low water conditions a higher proportion of the coho salmon present might have been caught than under high water conditions. Another possible explanation is that physical characteristics of the fish themselves (for example, sex or length) may influence both rate of capture for tagging and rate of recovery in tag recovery samples. For example, the beach seine may capture larger coho salmon at a higher rate than smaller coho salmon so that a higher proportion of the larger fish were tagged. As long as there is random mixing of coho salmon tagged during different time periods in the recovery areas, and the recovery process does not have the same selectivity as the capture process, this presents no problems for the abundance estimates.

Significant differences in the probability of finding a tag during surveys conducted at different times in a recovery area (p) were often found. Temporal trends in the physical characteristics of the population, combined with temporal trends in capture efficiency at the tagging site, could cause the changes observed. During spawning ground surveys, male fish may be more likely to end up in locations that are sampled than female fish, or larger fish may have a higher

probability of being seen and sampled during spawning ground surveys than smaller fish. The available data were examined to determine if these influences were present. The data used in these analyses were the length and sex composition data for all coho salmon tagged at the lower river tagging site and the tag recovery data used for the population estimates. Coho salmon recovered with a missing or illegible tag but having an operculum punch could not be used since their length and sex were not recorded at time of recovery.

Seber (1982) recommends testing the release (tagging) and recovery (escapement) samples for randomness with respect to length. The recovery sample was tested by comparing the length distributions of individuals that were tagged but not recovered to those individuals that were tagged and recovered. Both a Mann-Whitney U test and a Kolmogorov-Smirnov (K-S) test (Conover 1980) were used to compare the length distributions of coho salmon from these two groups. These same tests were also used to compare the length distributions of male and female coho salmon that were tagged in the lower Skagit River.

If there was a significant difference between the length distributions of male and female coho salmon subsequent analyses were conducted for each sex separately. If there was a significant difference between the length distributions of coho salmon which were tagged but not recovered and those that were tagged and recovered, K-S tests were performed sequentially on the length distributions to determine length categories with no significant difference between these two groups. Testing began between 65 and 70 cm (above which the length distributions of the two groups were not significantly different) and length was sequentially decreased by one cm intervals until a significant difference ($P \leq 0.05$) between the groups was found. A K-S test was then performed on those fish that were at the length of the significant difference or smaller. If there was a significant difference between the fish which were tagged but not recovered and those that were tagged and recovered the process was repeated for the fish in this smaller length range.

RESULTS

The results of the tagging conducted in 1988 are summarized in the following five sections. The summary consists of: (1) tag releases by day; (2) tag recoveries by location; (3) abundance estimates produced using the tag release-and-recovery data; (4) additional analyses which include migratory timing information from the release-and-recovery data and sex-length composition data; and (5) a discussion of the “best” estimate of the number of coho salmon migrating through the lower Skagit River tagging area.

There are two different tag recovery percentages presented in the results: the percentage of tags recovered from the tag releases during a specific time stratum (π) and the percentage of tagged fish in samples collected during tag recovery surveys (ρ). The recovery data from each major area were tested to determine if there were significant temporal differences in both of these percentages. The results of these tests determined which data were pooled and which model was used to estimate the abundance of coho salmon using the recovery data for a specific area or group of areas pooled.

Tag Releases

Tagging began on 12 September and continued through 15 November. A total of 1,970 coho salmon were tagged during 27 days of tagging (Table 1). About 24% of the tagged fish were eventually recovered during surveys conducted to estimate the percentage of tagged fish in the escapement.

The percentage of each day's release of tags that was recovered ranged from 0% to 50% (Figure 5). Generally, coho salmon tagged and released during September were recovered at a higher rate than those tagged and released in October and November. Four temporal release strata were defined to determine if there were significant differences in π among the release strata using the recoveries at each major area. The four release strata were:

1. 12 September to 20 September;
2. 23 September to 7 October;
3. 10 October to 21 October; and
4. 25 October to 15 November.

Significant differences in π among the release strata were found for the recoveries at Marblemount Hatchery, Suiattle sub-basin samples, the commercial fishery, and for the upriver spawning grounds combined (Table 2). There were no significant temporal differences in π among release strata at the other major recovery areas (Baker River trap, and Middle Skagit, Upper Skagit, and Lower Sauk sub-basin spawning grounds and for all recovery data combined). These tests were conducted only for recovery areas with seven or more legible tag recoveries.

Table 1. Number of coho salmon tagged each day and number of in-sample tag recoveries from each day's release for the Skagit River, 1988.

Date	Number Tagged	Tag Recoveries by Area ^a											Recoveries	
		MMH	BAK	MSK	USK	LSA	MSA	USA	SUI	OTH	CFS	TFS	Total	% (π)
12-Sep	34	1	0	0	0	0	0	0	0	0	1	0	2	5.9%
13-Sep	60	3	3	0	0	0	0	0	0	0	2	0	8	13.3%
15-Sep	22	0	0	0	0	0	0	0	0	0	0	0	0	0.0%
16-Sep	12	2	0	0	0	0	0	0	0	0	2	0	4	33.3%
19-Sep	45	5	1	0	0	0	0	1	0	0	4	0	11	24.4%
20-Sep	44	4	1	0	0	0	0	0	0	1	5	0	11	25.0%
23-Sep	64	5	0	0	0	0	0	0	0	0	11	0	16	25.0%
27-Sep	18	3	1	0	0	0	0	0	0	0	5	0	9	50.0%
28-Sep	35	1	4	0	0	0	0	0	0	0	8	0	13	37.1%
03-Oct	106	12	4	1	1	0	0	0	0	0	0	0	18	17.0%
04-Oct	61	8	4	0	0	1	0	0	0	0	0	0	13	21.3%
06-Oct	87	14	3	1	0	0	0	0	0	0	0	0	18	20.7%
07-Oct	164	26	6	1	0	0	0	0	0	0	0	1	34	20.7%
10-Oct	153	16	2	0	3	0	0	0	1	0	0	0	22	14.4%
11-Oct	52	6	5	0	0	0	1	0	0	0	0	0	12	23.1%
13-Oct	89	16	5	0	0	0	0	0	2	0	0	0	23	25.8%
14-Oct	295	42	12	4	2	2	1	0	1	3	0	0	67	22.7%
20-Oct	98	16	3	1	0	1	0	0	1	0	0	0	22	22.4%
21-Oct	176	20	8	1	1	0	0	0	0	0	0	0	30	17.0%
25-Oct	80	13	3	1	0	1	0	0	1	0	0	0	19	23.8%
26-Oct	66	3	2	0	0	0	0	0	0	0	1	0	6	9.1%
28-Oct	66	6	1	0	0	1	1	1	2	0	0	0	12	18.9%
01-Nov	37	2	1	0	0	1	1	0	1	0	1	1	8	21.6%
02-Nov	32	2	0	0	1	0	0	0	0	1	0	0	4	12.5%
09-Nov	18	2	0	0	1	0	0	0	0	0	1	0	4	22.2%
10-Nov	4	0	0	0	0	0	0	0	0	0	0	0	0	0.0%
15-Nov	52	3	0	0	0	0	0	0	0	0	4	0	7	13.5%
UNKNOWN ^b		32	8	5	2	7	6	2	5	3	1	0	71	
TOTALS	1,970	263	77	15	11	14	10	4	14	8	46	2	464	
% Recovered		13.4%	3.9%	0.8%	0.6%	0.7%	0.5%	0.2%	0.7%	0.4%	2.3%	0.1%	23.6%	

^a Locations are: MMH - Marblemount Hatchery; BAK - Baker River trap; MSK - Middle Skagit sub-basin; USK - Upper Skagit sub-basin; LSA - Lower Sauk sub-basin; MSA - Middle Sauk sub-basin; USA - Upper Sauk sub-basin; SUI - Suiattle sub-basin; OTH - Cascade, Nookachamps, and Carpenter sub-basins; CFS - Commercial fishery; and TFS - Test fishery.

^b Fish recovered with no tag but having the secondary mark (an operculum punch) or an illegible tag.




Figure 5. Percent of tags recovered during in-sample surveys from each day of release for coho salmon tagged in the Skagit River, 1988.

Table 2. Summary of the number of tag recoveries (#) from each release stratum in each major recovery area and the results of testing recovery percentages (π) for equality among release strata, 1988.

Release Strata		RECOVERY AREA											
		<u>Marblemount</u>		<u>Baker R.</u>		<u>Mid. Skagit</u>		<u>Up. Skagit</u>		<u>Lower Sauk</u>			
	Number Tagged	#	π	#	π	#	π	#	π	#	π	#	π
12-Sep thru 20-Sep	217	15	6.9%	5	2.3%	0	0.0%	0	0.0%	0	0.0%	0	0.0%
23-Sep thru 7-Oct	535	69	12.9%	22	4.1%	3	0.6%	1	0.2%	1	0.2%	1	0.2%
10-Oct thru 21-Oct	863	116	13.4%	35	4.1%	6	0.7%	6	0.7%	3	0.3%	3	0.3%
25-Oct thru 15-Nov	355	31	8.7%	7	2.0%	1	0.3%	2	0.6%	3	0.8%	3	0.8%
TOTALS	1,970	231	11.7%	69	3.5%	10	0.5%	9	0.5%	7	0.4%		
TEST RESULTS ^b		χ^2		χ^2		ART		ART		ART		ART	
Test Used:													
Significance (P):		0.01 ***		0.19 NS		0.46 NS		0.27 NS		0.29 NS			

- continued -

Table 2. Summary of the number of tag recoveries (#) from each release stratum in each major recovery area and the results of testing recovery percentages (π) for equality among release strata, 1988 (continued).

Release Strata	Number Tagged	RECOVERY AREA									
		<u>Suiattle</u>		<u>Spawn. Gr.^a</u>		<u>Comm. Fishery</u>		<u>Total</u>			
		#	π	#	π	#	π	#	π	#	π
12-Sep thru 20-Sep	217	0	0.0%	2	0.9%	13	6.0%	36	16.6%		
23-Sep thru 7-Oct	535	0	0.0%	5	0.9%	24	4.5%	121	22.6%		
10-Oct thru 21-Oct	863	5	0.6%	25	2.9%	0	0.0%	176	20.4%		
25-Oct thru 15-Nov	355	4	1.1%	14	3.9%	7	2.0%	60	16.9%		
TOTALS	1,970	9	0.5%	46	2.3%	44	2.3%	393	19.9%		
TEST RESULTS ^b											
Test Used:		ART								χ^2	
Significance (<i>P</i>):		0.04 ***		<0.01 ***		<0.01 ***		<0.01 ***		0.11 NS	

^a Total for all spawning ground samples from the Middle Skagit sub-basin and above. The total does not include recoveries from Marblemount Hatchery, Baker River trap, or the Nookachamps and Carpenter sub-basins.

^b Results of the tests to determine if the recovery percentages (π) were different among release strata. Test used: χ^2 = chi-square test, ART = approximate randomization test. NS = Not Significant; * = Significant, $0.05 < P \leq 0.10$; ** = Significant, $0.01 < P \leq 0.05$; *** = Significant, $P \leq 0.01$.

Tag Recoveries

Samples to estimate ρ were collected at 13 areas in the Skagit River drainage. A total of 24,552 coho salmon were examined of which 24,028 fish were considered in-sample and 524 were not considered part of the population subject to tagging. Sample surveys were conducted at: Marblemount Hatchery; Baker River trap; spawning grounds in the Middle Skagit, Upper Skagit, Lower Sauk, Middle Sauk, Upper Sauk, Suiattle, Cascade, Nookachamps, and Carpenter sub-basins; and in commercial and test fisheries. Of the 464 in-sample recoveries, 71 fish (15%) had a tag with an illegible number or had a missing tag and were identified as tagged by the opercula punches. Most of the in-sample recoveries were at Marblemount Hatchery (263 recoveries or 57% of all in-sample recoveries). The areas with the next largest number of tag recoveries were Baker River (77 or 17%) and the commercial fishery (45 or 10%). Combined, these three areas account for 83% of all in-sample recoveries.

The percentage of tagged fish in the escapement samples (ρ) from the nine recovery areas having seven or more tag recoveries ranged from 1.2% for Middle Skagit sub-basin samples to 3.7% for Cascade sub-basin samples (Table 3). There was a significant difference (χ^2 , $P < 0.01$) in ρ among these nine areas.

The average number of days between release and recovery for in-sample tag recoveries was about 39 days (Table 4). Tagged coho salmon recovered in the test fishery had the shortest average time between release and recovery, 7 days, and tag recoveries from the Middle Sauk sub-basin had the longest average time between release and recovery, 90 days. For the upstream recovery areas, tag recoveries at Baker River trap had the earliest average day of release (10 October) and recoveries from the Middle Sauk sub-basin had the latest average day of release (21 October).

Marblemount Hatchery:

Escapement samples were collected at Marblemount Hatchery from 19 October through 17 January. A total of 8,783 coho salmon were examined and 263 tagged fish (3.0%) were found (Appendix Table A-1). The Marblemount Hatchery sample is considered a census because all returning fish are sampled so the data were not examined for temporal differences in ρ .

Baker River Trap:

Escapement samples were collected at Baker River trap from 9 September through 27 January. A total of 2,820 coho salmon were examined for tags. Based upon a four-day minimum travel time from the tagging area to Baker River dam determined from all five years of tagging data (Conrad et al. 1997), samples collected prior to 16 September were not considered in-sample since tagging did not begin until 12 September. A total of 2,781 coho salmon were examined for tags from 16 September through 27 January and 77 tagged fish (2.8%) were found (Appendix Table A-2). The Baker River trap sample is considered a census because all returning fish are sampled so the data were not examined for temporal differences in ρ .

Table 3. Summary of the percentage of tagged or marked coho salmon found in each recovery area during in-sample surveys of the Skagit River, 1988.

Recovery Area	Time Period	Fish Examined	Tags Found ^a	% Tagged (p)
Marblemount Hatchery	1. 19-Oct - 17-Jan	8,783	263	3.0%
Baker River Trap	X. ^b 9-Sep - 13-Sep	39	0	0.0%
	1. 16-Sep - 27-Jan	2,781	77	2.8%
Commercial Fishery	1. 22-Sep - 30-Sep	1,194	37	3.1%
	2. 13-Dec - 20-Jan	934	8	0.9%
	Total	2,128	45	2.1%
Middle Skagit Sub-basin	1. 20-Oct - 23-Feb	848	5	0.6%
Etach Slough Trap	1. 01-Nov - 30-Nov	122	7	5.7%
	2. 01-Dec - 11-Feb	305	3	1.0%
	Total	1,275	15	1.2%
Lower Sauk Sub-basin	1. 03-Nov - 27-Feb	654	14	2.1%
Suiattle Sub-basin	1. 14-Nov - 14-Feb	679	14	2.1%
Upper Skagit Sub-basin	1. 27-Oct - 29-Mar	479	11	2.3%
Middle Sauk Sub-basin	1. 22-Oct - 23-Feb	683	10	1.5%
Cascade Sub-basin	1. 07-Nov - 23-Feb	191	7	3.7%
Upper Sauk Sub-basin	1. 02-Nov - 15-Feb	50	4	8.0%
IN-SAMPLE TOTAL FOR UPSTREAM AREAS		17,703	460	2.6%
Carpenter Sub-basin	1. 23-Oct - 27-Jan	838	0	0.0%
Nookachamps Sub-basin	1. 04-Nov - 13-Jan	419	1	0.2%
Commercial Fishery (downstream)	X. 9-Sep	154	0	0.0%
	1. 13-Sep - 19-Jan	2,792	1	<0.1%
Test Fishery (downstream)	X. 17-Aug - 7-Sep	331	0	0.0%
	1. 14-Sep - 04-Nov	2,276	2	0.1%
IN-SAMPLE TOTAL FOR DOWNSTREAM AREAS		6,325	4	0.1%
TOTAL CONSIDERED IN POPULATION BEFORE TAGGING		524	0	0.0%
IN-SAMPLE TOTAL FOR ALL AREAS		24,028	464	1.9%
GRAND TOTAL FOR ALL SAMPLES		24,552	464	1.9%

^a Includes fish recovered with no tag but having the secondary mark (an operculum punch) or having an illegible tag.

^b X indicates that these fish were considered to be in the population before tagging began and not subject to tagging (i.e., they were not considered in-sample fish for the abundance estimates).

Table 4. Average day of release (DOR) and average number of days between release and recovery (DBET) for coho salmon tagged and recovered in the Skagit River, 1988.

Recovery Area	Sample Size ^a	Average DOR	Stand. Error	Range	Average DBET	Stand. Error	Range
Marblemount Hatchery:							
Pond Mortalities	17	11-Oct	1.9	27-Sep - 21-Oct	62.2	3.4	40 - 88
Surplussed	17	25-Sep	2.5	16-Sep - 28-Oct	28.5	2.8	15 - 68
Spawned	197	13-Oct	0.7	12-Sep - 15-Nov	42.3	0.9	7 - 79
Baker River Trap	67	10-Oct	1.3	13-Sep - 1-Nov	21.4	1.6	5 - 85
Commercial Fishery	45	30-Sep	2.7	12-Sep - 15-Nov	11.1	2.0	1 - 50
Middle Skagit Sub-basin	10	13-Oct	2.2	3-Oct - 25-Oct	53.6	7.0	27 - 101
Lower Sauk Sub-basin	7	19-Oct	3.7	4-Oct - 1-Nov	78.9	7.8	55 - 116
Suiattle Sub-basin	9	20-Oct	2.7	10-Oct - 1-Nov	78.8	5.6	47 - 102
Upper Skagit Sub-basin	9	17-Oct	4.0	3-Oct - 9-Nov	73.3	9.9	26 - 119
Middle Sauk Sub-basin	4	21-Oct	5.2	11-Oct - 1-Nov	90.0	12.6	66 - 125
Cascade Sub-basin	5	13-Oct	6.8	20-Sep - 2-Nov	79.8	8.2	47 - 91
Upper Sauk Sub-basin	2	8-Oct	19.5	19-Sep - 28-Oct	74.5	30.5	44 - 105
Test Fishery	2	19-Oct	12.5	7-Oct - 1-Nov	7.0	5.0	2 - 12
All Recoveries	391	10-Oct	0.6	12-Sep - 15-Nov	38.8	1.2	1 - 125

^a Includes tag recoveries with legible numbers only.

Commercial and Test Fishery Samples:

A commercial fishery was conducted in the river and in Skagit Bay on 37 days between 9 September and 20 January. Catches from areas above and below the tagging site were sampled. Recovery data collected on 9 September were excluded from analysis since tagging did not begin until 12 September. A total of 2,128 coho salmon were examined for tags and 45 tagged fish (2.1%) were found in catches from areas above the tagging site (Appendix Table A-3). For the upstream samples, there was a significant difference (χ^2 , $P < 0.01$) in p between the catches sampled in September and those sampled in December and January. After tagging began, only one tagged coho salmon was found in 2,792 fish examined for tags from catches in downstream areas (including Skagit Bay). Recovery data collected from the downstream areas were not used for the population estimates.

Test fisheries were conducted on 24 days between 17 August and 4 November. Recovery data collected from 17 August through 7 September were excluded from analysis since tagging did not begin until 12 September. A total of 2,276 coho salmon were examined for tags after tagging began and only two tagged fish (0.1%) were found (Appendix Table A-4).

Middle Skagit Sub-basin:

Tag recovery samples were collected during surveys of Middle Skagit sub-basin spawning grounds conducted from 20 October through 23 February and at two traps on tributaries to the sub-basin. Surveys were conducted by SSC crews. A total of 848 coho salmon were examined for tags and 5 tagged fish (0.6%) were found during the surveys and at the trap on Careys Creek (Appendix Table A-5). The percentage of tags in samples collected from the trap on Etach Slough was significantly different (χ^2 , $P < 0.01$) from the percentage in samples collected during spawning ground surveys and at the other trap, therefore, the data were not combined. There was not a significant difference in p between samples collected on the rack and at the trap on Etach Slough (Fisher's exact test, $P = 0.13$) so these samples were combined. A total of 427 coho salmon were examined for tags and 10 tagged fish (2.3%) were found at the trap on Etach Slough (Appendix Table A-6). There was a significant difference (Fisher's exact test, $P < 0.01$) in p between samples collected from 1 November through 30 November and samples collected after 30 November.

Upper Skagit Sub-basin:

Tag recovery samples were collected during surveys of Upper Skagit sub-basin spawning grounds conducted from 27 October through 29 March. Surveys were conducted by SSC crews. A total of 479 coho salmon were examined for tags and 11 tagged fish (2.3%) were found (Appendix Table A-7). The hypothesis of constant p for temporal strata in the recovery samples could not be rejected.

Lower Sauk Sub-basin:

Tag recovery samples were collected during surveys of Lower Sauk sub-basin spawning grounds conducted from 3 November through 27 February. Surveys were conducted by SSC crews. A total of 654 coho salmon were examined for tags and 14 tagged fish (2.1%) were found (Appendix Table A-8). The hypothesis of constant p for temporal strata in the recovery samples could not be rejected.

Middle Sauk Sub-basin:

Tag recovery samples were collected during surveys of Middle Sauk sub-basin spawning grounds conducted from 22 October through 23 February. Surveys were conducted by SSC crews. A total of 683 coho salmon were examined for tags and 10 tagged fish (1.5%) were found (Appendix Table A-9). The hypothesis of constant p for temporal strata in the recovery samples could not be rejected.

Upper Sauk Sub-basin:

Tag recovery samples were collected during surveys of Upper Sauk sub-basin spawning grounds conducted from 2 November through 15 February. Surveys were conducted by SSC crews. Only 50 coho salmon were examined for tags and four tagged fish (8.0%) were found (Appendix Table A-10).

Suiattle Sub-basin:

Tag recovery samples were collected during surveys of Suiattle sub-basin spawning grounds conducted from 14 November through 14 February and at a trap on Suiattle Slough operated by WDFW from 26 November through 6 February. Surveys were conducted by SSC crews. There was not a significant difference in p between samples collected during surveys and at the trap (Fisher's exact test, $P = 0.37$) so the samples were combined. A total of 679 coho salmon were examined for tags and 14 tagged fish (2.1%) were found (Appendix Table A-11). The hypothesis of constant p for temporal strata in the recovery samples could not be rejected.

Other Spawning Ground Surveys:

Spawning ground surveys were conducted in three other areas: Nookachamps sub-basin, Carpenter sub-basin, and Cascade sub-basin. Tag recovery samples were collected during surveys of Nookachamps sub-basin spawning grounds by SSC and at traps on the East and West Forks of Nookachamps Creek. A total of 419 coho salmon were examined for tags and only one tagged fish (0.2%) was found (Appendix Table A-12). Spawning ground surveys of the Carpenter sub-basin were conducted by SSC crews and a trap was operated by SSC on Carpenter Creek Slough. A total of 838 coho salmon were examined for tags but no tagged fish (0.0%) were found in these samples (Appendix Table A-13). SSC crews surveyed Cascade sub-basin spawning grounds and examined 191 coho salmon (Appendix Table A-14). Seven tags were recovered during surveys of the Cascade sub-basin (3.7%).

Out-of-System Recoveries:

There were no jaw tags from the tagging conducted in the Skagit River during 1988 that were recovered outside of the Skagit River system.

Abundance Estimates

Estimates of coho salmon abundance from the tag recovery data for each major recovery area having seven or more tag recoveries are summarized in Table 5. The details of the abundance estimate for each location are in Appendix B. The commercial fishery samples from the upstream areas were not used to estimate abundance because the fishery occurred during two discrete time periods (with a 73 day interval between them) and p for these two periods was significantly different (χ^2 , $P < 0.01$). There was also a significant difference in p between the samples collected at Etach Slough and at other Middle Skagit sub-basin locations, and there was a significant difference in p between temporal strata at Etach Slough. There was not a logical way to combine the data from the two temporal strata in Etach Slough with the other Middle Skagit sub-basin data which overlapped both of the time strata, so no estimate was generated using Middle Skagit sub-basin recovery data. The samples from Marblemount Hatchery and Baker River trap were both censuses so they were compared to determine if it was appropriate to pool them. The two samples were not significantly different (χ^2 , $P = 0.54$) so an estimate was generated for the pooled data.

The five samples from sub-basins above the tagging area which had no significant temporal differences in p (Lower Sauk, Middle Sauk, Suiattle, Upper Skagit, and Cascade) were compared and no significant differences in p were found (χ^2 , $P = 0.44$). The recovery data from these locations were pooled to generate an estimate. Finally, p for Marblemount Hatchery, Baker River trap, and the Lower Sauk, Middle Sauk, Suiattle, Upper Skagit, and Cascade sub-basins were compared. The hypothesis of equal p was not rejected (χ^2 , $P = 0.17$) so these samples were pooled for an estimate.

Estimates of the number of coho salmon migrating through the tagging area in the lower Skagit River ranged from 47,303 coho salmon using Cascade sub-basin spawning ground recovery data to 122,559 coho salmon using Middle Sauk sub-basin spawning ground recovery data. Pooled Marblemount-Baker-Lower Sauk-Suiattle-Upper Skagit-Middle Sauk-Cascade data provided the most precise estimate (CV = 4.4%). The estimate with the largest CV was from Cascade sub-basin recovery data (CV = 32.6%). The 95% confidence intervals for the abundance estimates overlapped for each recovery area.

Table 5. Summary of estimates of the number of coho salmon in the Skagit River escapement using data from each major recovery area, 1988.

Recovery Area	Estimation Method	Estimated Abundance	Stand. Error	CV ^a	95% Confidence Interval
Marblemount	Petersen	65,580	3,692	5.6%	58,790 - 74,679
Baker River	Petersen	70,298	7,642	10.9%	58,306 - 91,252
Marblemount - Baker pooled	Petersen	66,845	3,238	4.8%	60,651 - 74,841
Lower Sauk	Petersen	86,066	21,188	24.6%	49,216 - 155,894
Suiattle	Petersen	89,351	22,006	24.6%	51,097 - 161,853
Upper Skagit	Petersen	78,839	21,525	27.3%	41,803 - 155,699
Middle Sauk	Petersen	122,559	34,996	28.6%	62,970 - 252,956
Cascade	Petersen	47,303	15,405	32.6%	21,109 - 116,267
L. Sauk - Suiattle - U. Skagit - M. Sauk - Cascade pooled	Petersen	92,913	11,894	12.8%	75,038 - 127,554
Marblemount - Baker - L. Sauk - Suiattle - U. Skagit - M. Sauk - Cascade pooled	Petersen	70,751	3,125	4.4%	64,614 - 78,516

^a CV = coefficient of variation.

Additional Analyses

The release data were divided into ten, 10-day time periods for the migratory timing analysis and to describe temporal patterns in the length and sex composition of tagged coho salmon. Coho salmon were tagged and released during seven of these periods.

Timing of Migrations to Major Recovery Areas:

The CPUE of coho salmon by the beach seine in the lower river tagging area is shown by day and for each 10-day period in Figure 6. CPUE peaked during the 11 October through 20 October time period. Three areas had ten or more recoveries of legible tags which could be used for the migratory timing calculations (Appendix Table A-15). The trends in CPUE of Marblemount Hatchery fish and fish bound for Baker River were very similar to the trend for total CPUE by 10-day period (Figure 7). Coho salmon from every period with tag releases were recovered at Marblemount Hatchery: this was the only area in which this occurred.

Length and Sex Composition Analyses:

The sex and length data for the 1,970 coho salmon tagged and released in the lower Skagit River and the 391 in-sample recoveries with legible tags were analyzed. Both the K-S and M-W tests which compared the lengths of coho salmon tagged but not recovered to the lengths of those tagged and recovered were significant ($P < 0.01$) indicating that the recovery samples were not random with respect to length of fish. There was also a significant difference between male and female length distributions (K-S test, $P < 0.01$), therefore, all subsequent analyses of length were conducted for each sex separately. It is evident from Figure 8 that male coho salmon had a higher proportion of smaller sizes (fish less than 50 cm) than female coho salmon. Coho salmon less than 50 cm in length composed about 36% of the males that were tagged but only 4% of the female coho salmon that were tagged.

Males Tagged male coho salmon averaged 53.1 cm in fork length (SE = 0.24). The mean length of male coho salmon that were tagged but not recovered was 52.5 cm (SE = 0.27) compared to a mean length of 55.5 cm (SE = 0.44) for male coho salmon that were tagged and recovered. The length distribution of male coho salmon that were tagged but not recovered was significantly different (K-S test, $P < 0.01$) from the distribution of those that were tagged and recovered (Figure 9). Two length categories were defined from the sequential K-S tests for male coho salmon: (1) fish with lengths less than 50 cm; and (2) fish with lengths equal to or greater than 50 cm. The percentages of tagged coho salmon in each length category that were recovered were 9.7% and 25.9%, respectively (Appendix Table A-16).




Figure 6. Catch-per-unit effort of coho salmon by the beach seine in the lower Skagit River tagging area by day and for each ten-day period, 1988.




Figure 7. Beach seine catch-per-unit effort (CPUE) of coho salmon bound for major Skagit River tag recovery areas in 1988. CPUE is for ten-day periods (starting date of period shown) and is expressed as a percentage of the total CPUE for tagged fish recovered from the area.




Figure 8. Comparison of length frequencies of male and female coho salmon tagged in the lower Skagit River, 1988.




Figure 9. Comparison of length frequencies of coho salmon that were tagged but not recovered to those that were tagged and recovered, for males and females, 1988.

Females Tagged female coho salmon averaged 58.4 cm in fork length (SE = 0.19). The mean length of female coho salmon that were tagged but not recovered was 58.6 cm (SE = 0.21) compared to a mean length of 57.4 cm (SE = 0.39) for female coho salmon that were tagged and recovered. The length distribution of female coho salmon that were tagged but not recovered was significantly different (K-S test, $P < 0.01$) from the distribution of those that were tagged and recovered (Figure 9). Two length categories were defined from the sequential K-S tests for female coho salmon: (1) fish with lengths less than 59 cm; and (2) fish with lengths equal to or greater than 59 cm. The percentages of tagged coho salmon in each length category that were recovered were 24.0% and 14.2%, respectively (Appendix Table A-16).

Tag Recovery Rates There was not a significant difference (χ^2 , $P = 0.71$) in tag recovery rates between male and female coho salmon. The highest rate of tag recovery, 25.9%, was for males in the largest length category (≥ 50 cm) and the lowest tag recovery rate (9.7%) was for males in the smallest (≤ 49 cm) length category (Appendix Table A-16). There was not a significant difference (χ^2 , $P = 0.18$) in tag recovery rates among the release condition categories. There was a significant difference (χ^2 , $P = 0.03$) in tag recovery rates among the maturity categories. Coho salmon classified as bright had a 19.1% tag recovery rate while those classified as dark had a 35.7% tag recovery rate (Appendix Table A-17).

Sex-Length Composition There were temporal changes in both the sex composition and length composition for each sex during the tagging period (Figure 10). The percentage of males in the tagging samples gradually declined (from 61% to 49%) throughout the release period and the percentage of females gradually increased (from 39% to 51%). The percentage of small males (≤ 49 cm) decreased throughout the release period as the percentage of large males (≥ 50 cm) increased. The percentage of small and large females started at 78% and 22%, respectively, and gradually converged to about equal proportions by the end of the tagging period.

Conclusions

The percentage of tagged coho salmon (p) in the samples from the major recovery areas (areas with seven or more tag recoveries) was not as consistent as in 1986 and 1987. About 3% of the coho salmon returning to Marblemount Hatchery and the Baker River trap were tagged, while p for most of the samples from the sub-basin spawning grounds above the tagging area was about 2% (Lower Sauk sub-basin 2.1%; Suiattle sub-basin 2.1%; Upper Skagit sub-basin 2.3%), but these differences were not significant. The exceptions were the Middle Skagit and Middle Sauk sub-basins where 1.2% and 1.5%, respectively, of the fish examined were tagged or marked and the Cascade sub-basin where p was 3.7%. The tag recovery data indicate that some coho salmon from spawning areas substantially downstream of the tagging site were present in the tagging area. There was one tag recovered in 1,257 coho salmon examined (0.08%) during spawning ground surveys in the Nookachamps and Carpenter sub-basins (Figure 1).




Figure 10. Sex and length composition, by release period, of coho salmon tagged in the lower Skagit River, 1988.

We recommend that the estimate using the pooled Marblemount-Baker-Lower Sauk- Suiattle-Upper Skagit-Middle Sauk-Cascade data be considered the “best” estimate of coho salmon abundance for 1988. There was not a significant difference in ρ among these areas. The samples from two of the areas, Marblemount Hatchery and Baker River trap, are censuses. Combined, these two areas accounted for 86% of the tag recoveries and 81% of the fish examined in the data used for the population estimate. There were no temporal differences in ρ in the samples from the five sub-basins. This estimate uses the largest number of tag recoveries (396) and therefore has the smallest CV. The estimate, 70,751 coho salmon (95% confidence interval: 64,614 to 78,516), is for the number of fish present in the lower Skagit River tagging area during the period 12 September to 15 November. The estimate includes coho salmon bound for all spawning grounds above the tagging area and some portion of the escapement to areas downstream of the tagging site. We do not recommend combining the recovery data from the commercial fishery or the Middle Skagit sub-basin because of the temporal differences in ρ for the samples from these recovery areas.

DISCUSSION

The number of coho salmon in the escapement to the Skagit River was estimated using the tag release-and-recovery data and the Petersen model. A discussion of how well the data meet the major assumptions of the Petersen model and a definition of the “population” which is being estimated follows.

Population was Closed

We assume that some coho salmon migrated through the tagging area before and after the period of tagging (12 September through 15 November). Although the Petersen model generally assumes a closed population, the population can be open but the exact point in time to which the estimate applies must be specified (Seber 1982). We feel the trend in CPUE for the beach seine used to capture coho salmon for tagging provides strong evidence that the tagging period encompassed the major portion of the coho salmon migration. The CPUE was low when tagging began and was followed by an increase in CPUE to a peak during the period 11 October through 20 October. This was followed by a decline in CPUE in late October and early November (Figure 6).

Similarly to 1986 and 1987, adjustments were made to the total number of fish examined at Baker River trap to account for early-arriving fish that were not subject to tagging. Therefore, the estimate includes only the portion of the population migrating through the tagging area after tagging began.

If we assume there is recruitment to the population (coho salmon migrating through the tagging area after tagging ends) but no mortality before the fish reach their spawning areas, and there is complete mixing of the fish on the spawning grounds, then the abundance estimate includes coho salmon migrating through the tagging area after the last day of tagging. Sampling at Marblemount Hatchery and at Baker River trap was conducted through 17 and 27 January, respectively. Tag recovery surveys were conducted in most sub-basin spawning grounds until mid or late February. We feel there was sufficient time for coho salmon migrating through the tagging area after tagging had ended to mix with the fish already present on the spawning grounds and at Marblemount Hatchery.

Area Encompassed by the Estimates:

The Petersen model estimates the number of coho salmon migrating through the tagging area in the lower river during the time period defined above. The estimate includes all coho salmon bound for spawning areas above the tagging area (including Marblemount Hatchery and Baker River) and all spawning areas in the Middle Skagit sub-basin above and including Hansen Creek (Figure 1). However, the tag release-and-recovery data suggest that only a fraction of the coho salmon which spawned in the Carpenter and Nookachamps sub-basins passed through the tagging area. The percentage of tags in the samples from these areas combined, 0.06% (including downstream commercial and test fishery samples), was much smaller than in the

upstream recovery areas. This indicates that only a portion of the coho salmon from these areas passed through the tagging area. Therefore, we conclude that the abundance estimate does not include all of the coho salmon which spawned in the Carpenter and Nookachamps sub-basins. If the total number of tagged fish that migrated to these downstream areas could be estimated, this number could be removed from the total number of tags released and the abundance estimate would include only coho salmon bound for areas **upstream** of the tagging site and the Middle Skagit sub-basin. We estimated the number of tags “lost” to these downstream areas so that we could examine the effect of these tags on the abundance estimate for the upstream areas.

Estimate of the Number of Tagged Fish “Lost” to Areas Downstream of the Tagging Area:

Tagged coho salmon were present in three groups of fish from areas downstream of the tagging area: (1) commercial fishery catches; (2) test fishery catches, and (3) fish spawning in the Carpenter and Nookachamps sub-basins. The commercial catch in area 78D was sampled by subareas (78D-2, 78D-3, and 78D-4; see Figure 4) in 1988 so we could estimate the percentage of tags in samples above and below the tagging area. The total commercial catch from area 78D is not recorded by these sub-areas, however. Therefore, we assumed that the proportional catch by sub-area for the commercial catch samples was the same as the entire commercial catch for area 78D. We pooled all catches and applied the percentage of tags found in downstream commercial and test fishery samples (Area 2, Spudhouse, Blakes, Bay and Jetty; see Figure 4) to the pooled total. The number of tagged fish present on spawning grounds in the Carpenter and Nookachamps sub-basins was estimated by applying the percentage of tags found during in-sample surveys of these sub-basins combined (one tagged fish found in 1,257 fish examined for $p = 0.08\%$) to an independent estimate of the number of coho salmon spawning in these sub-basins. The spawning ground escapement to these sub-basins was estimated using a redd-count method (Conrad et al. 1993). The numbers used for these calculations are summarized in Appendix Table A-18. We estimated that a total of 10 tags could have been “lost” to these downstream areas. If the number of tags released is adjusted to 1,960 (1,970 - 10), then (using the pooled Marblemount-Baker-Lower Sauk-Suiattle-Upper Skagit-Middle Sauk-Cascade recovery data) the estimated abundance for areas upstream of the tagging area becomes 70,392 coho salmon. This is only 359 fish less than the “unadjusted” estimate and is less than a one percent difference.

The presence of coho salmon in the tagging area bound for systems outside the Skagit River would also affect the abundance estimate. In 1988, there were no out-of-system recoveries of coho salmon tagged in the lower Skagit River. Therefore, we do not feel that the contribution of coho salmon bound for systems outside the Skagit River was a major source of error.

All Coho Salmon Have an Equal Probability of Capture During Tagging or the Recovery Sample is a Simple Random Sample of the Population

These assumptions are often hard to satisfy as it is difficult or impossible to obtain simple random samples from highly dispersed and mobile populations. Fortunately, the estimates are still valid under certain alternative assumptions. Junge (1963) demonstrated that selectivity (non-randomness) may exist in both the tagging and recovery samples without introducing bias in the estimate if the sources of selectivity in the two samples are independent.

During the Skagit River study, there is evidence that the tagging sample may not have been random with respect to time. Certain portions of the population may have been tagged at higher rates than others. There is also evidence that the recovery samples on the spawning grounds were selective with respect to the length of the fish, at least for males. Eames et al. (1981, 1983) found that there was a correlation between time of entry and size of coho salmon for the returns to the Skagit River in 1976 and 1977. Smaller fish generally arrived earlier in the run than larger fish. This presents a problem if timing of passage through the tagging area is correlated with the size of fish and area of spawning (Junge 1963). If such selectivity existed the population estimates would contain a negative bias. However, we believe if such a bias exists it is small because the majority of the tag recovery data used for the abundance estimate was collected from areas where there was no size selectivity (Marblemount Hatchery and Baker River trap).

The use of different gears to obtain the tagging and recovery samples is a common technique for minimizing the bias due to selectivity (Ricker 1975; Seber 1982). In this study, coho salmon were captured for tagging using a beach seine. Recovery samples were either a census of all adults returning to an area (Marblemount Hatchery and Baker River trap) and thus non-selective, or were samples collected on the spawning grounds during foot surveys (and to a lesser extent by traps in some areas). We do not feel that selectivity (non-random sampling) was a significant source of bias for the estimates because: (1) the methods used to capture coho salmon for tagging were different from those used to recover them; and (2) a majority of the tag recoveries used to estimate abundance were collected by a census.

Tagging Does Not Affect the Catchability of an Animal

This assumption is necessary because some of the coho salmon passing through the tagging area were subject to an in-river commercial fishery above the tagging area. If jaw-tagged coho salmon were removed at a different rate than untagged fish, the percentage of tags in any recovery samples collected after this removal would be different from the percentage of tags in the population immediately after tagging. There is no evidence of selective removal of tagged fish in the data. In 1988, the percentage of tagged fish in the commercial fishery samples from sub-areas of 78D upstream of the tagging area was essentially the same as that observed on most upstream spawning grounds with seven or more tag recoveries (with the exception of the Cascade sub-basin). The percentage of tags in the commercial fishery samples was less than that observed at Marblemount Hatchery and Baker River trap. If tags were being removed

selectively, we would expect the commercial fishery samples to have a higher percentage of tags than these samples.

Animals Do Not Lose Their Tags Between the First and Second Samples

In 1988, 15% of the tagged coho salmon recovered had missing or illegible tags. However, the use of opercula punches on all tagged fish allowed coho salmon with missing tags to be identified as previously tagged. Identified tag loss must be accounted for only in the Darroch estimate of abundance which requires that the release period of recovered individuals be known. When there was no tag but an operculum punch was present (or the tag was illegible), the release period was estimated as described in the Methods section. This was required only when the Darroch estimate was selected as the appropriate model. The Darroch estimate was not used for any of the abundance estimates produced in 1988. The Petersen estimate was selected as the appropriate model for all estimates. As long as all coho salmon with a missing tag are identified by an operculum punch, the Petersen estimate is not affected by the missing tags.

All Tagged Animals are Reported in the Second Sample

Because the majority of the tag recoveries used for the abundance estimates were from Marblemount Hatchery, and all coho salmon at Marblemount Hatchery were inspected twice for tags, we expect very few jaw-tagged (or marked) fish were missed. Live fish were individually inspected for tags and marks at Baker River dam. During surveys of spawning grounds, surveyors carefully inspected each carcass for an operculum punch if no tag was visible. Considering that some carcasses were in an advanced state of decay it is possible that some fish with a missing tag were not identified. In 1988, about 6% of the carcasses examined on the upriver spawning grounds (Middle Skagit sub-basin and above) could not be sampled because of their condition.

There are No Mortalities Due to Tagging

Tests to determine the extent of tagging mortality were conducted during four of the five study years. These tests and their results are documented in Conrad et al. (1997). Based on these tests we concluded that there was no evidence of tagging mortality. The tests provided strong evidence that there was no short-term (within 48 hours) tagging mortality. The tag recovery data from the commercial fishery samples provide additional evidence that there was no delayed tagging-induced mortality occurring from two weeks up to three months after tagging. The average time between tag release and recovery for the commercial fishery recoveries, about 11 days (Table 4), was the shortest of any of the upstream recovery areas. Since the coho salmon caught in the commercial fishery are caught relatively soon after tagging, we would expect that if there is any delayed mortality caused by tagging it would cause the commercial fishery samples to have a higher percentage of tags than the samples that are taken much later, further upstream. In 1988, p for the commercial fishery samples from upstream areas (2.1%) was very similar to that for most of the upstream recovery areas.

CONCLUSIONS

The estimated abundance of coho salmon in 1988 was 70,751 fish with a 95% confidence interval of 64,614 to 78,516 fish. The mark-recapture estimate is for the number of coho salmon migrating through the tagging area after tagging began on 12 September. It includes all coho salmon bound for spawning areas above the tagging area and an unknown fraction of the salmon from spawning areas in the Nookachamps and Carpenter sub-basins. This abundance estimate was very precise ($CV = 4.4\%$) because of the large number of fish examined for tags during in-sample surveys. To restrict the estimate to spawning areas in the Middle Skagit sub-basin and spawning areas above it, adjustments were made to the number of tags released. Using the adjusted number of tags released, the estimated abundance for this more restricted area was 70,392 coho salmon. The variance of this estimate was not calculated because of the unknown precision for the estimated number of tags “lost” to downstream areas. The adjusted estimate falls within the 95% confidence interval of the original estimate.

To estimate the number of “wild” coho salmon which reached upstream spawning areas in the Skagit River during 1988, the number of hatchery fish plus any catches by the commercial and test fisheries above the tagging area need to be removed from the adjusted estimate and the number of fish which migrated through the tagging area prior to tagging needs to be added. Since fish which migrated through the tagging area before tagging began are included in the spawning ground samples, only prior-migrating fish returning to Baker River or caught in the commercial fishery need to be included. Since the return to the Baker River trap was censused, we have a total count of the prior-migrating fish to this area, 39 fish. Although all in-river catches were recorded on fish tickets, the sub-area of harvest within area 78D was not indicated on the ticket. Therefore, it was necessary to estimate the distribution of the commercial catch in area 78D to areas above and below the tagging site using the distribution of the commercial catch samples among areas 78D-1, 78D-2, 78D-3, and 78D-4. In-population sport catches should also be subtracted from the adjusted estimate. In-river catches of coho salmon by the sport fishery in the Skagit River were estimated to be only 1,050 fish in 1988 (WDF 1992) and were not included in the summary total as the specific dates and areas of harvest of these fish are unknown. A summary of the total terminal area run of coho salmon to the Skagit River in 1988 is presented in Table 6. **The total terminal area run of coho salmon to the Skagit River in 1988 is estimated to be 86,265 fish. An estimated 61,283 coho salmon were in the “wild” escapement to Skagit River spawning grounds:** 58,029 fish were estimated to have reached upstream spawning grounds and 3,254 coho salmon were estimated for lower river (Nookachamps and Carpenter sub-basin) spawning grounds. For comparison, the escapement of “wild” coho salmon to Skagit River spawning grounds estimated using index area surveys was 19,000 fish (Jeff Parkhurst, WDFW, personal communication). An alternative estimate, derived from CWT recoveries in the test fisheries and trap recoveries (Hayman 1996), was for a wild escapement of 64,000 to 72,000 fish (depending upon the hatchery stray rate assumed); this estimate was subsequently refined for a wild escapement estimate of 67,600 fish (Hayman 1997). Using a redd-count method, Conrad et al. (1993) estimated the wild escapement to be 75,000 to 112,000 fish (assuming two or three coho salmon per redd, respectively).

Table 6. Summary of the number of coho salmon returning to Skagit Bay in 1988.

Component	In-Population	Out of Population	Total
Upstream Estimated Total	70,392	40	70,432
Marblemount Hatchery	8,783	0	8,783
Baker River Hatchery ^a	1,499	1	1,500
Area 78D-3, 78D-4 Commercial Catch	2,119	1	2,120
Upstream Test Fishery Catch	0	0	0
Upstream Removals and Hatchery Fish	12,401	2	12,403
Estimated “Wild” Escapement to Upstream Spawning Areas	57,991	38	58,029
Nookachamps Sub-basin Estimated Escapement		1,405	1,405
Carpenter Sub-basin Estimated Escapement		1,849	1,849
Areas 78D-1, 78D-2, 78C, 8E, 8 Commercial Catches		9,959	9,959
Downstream Test Fishery Catch		2,620	2,620
Downstream Total		15,833	15,833
“Wild” Escapement ^b to Spawning Grounds	57,991	3,292	61,283 ^c
Total Terminal Run to Skagit Bay	70,392	15,873	86,265 ^c

^a Total number of hatchery coho salmon that returned to the Baker River trap. All coho salmon smolts from the Baker River Hatchery in the 1985 brood year (which primarily returned during 1988) were adipose fin clipped. The total return of coho salmon to the Baker River trap was 2,820 fish in 1988. Of these, 1,500 fish had adipose fin clips (i.e., were hatchery fish) and 1,320 were unmarked fish (assumed to be wild coho salmon). Of the unmarked fish, 38 returned prior to the first day of tagging and were considered out-of-population. The wild totals are included in the “wild” escapement numbers.

^b Includes estimated “wild” escapement to upstream spawning areas and estimated escapement to the Nookachamps and Carpenter sub-basins (from Conrad et. al 1993).

^c The estimated catch by the in-river sport fishery was 1,050 coho salmon, but the specific dates and areas of harvest of these fish are unknown. The total wild escapement should be reduced by the number of coho salmon caught in the sport fishery in upstream areas after tagging began. The total terminal run should be increased by the number caught in downstream areas or before tagging started.

ACKNOWLEDGMENTS

Funding for this project was administered by the Northwest Indian Fisheries Commission (NWIFC) under contract number #31000-643. Funding was provided pursuant to a PL-638 contract between the NWIFC and the U. S. Department of the Interior to meet obligations of the United States under terms of the Pacific Salmon Treaty.

Many people contributed in various ways to this study and it is not possible to list them all. However, we would like to acknowledge the contributions of the following groups of people: Washington Department of Fish and Wildlife: Marblemount Hatchery staff and Mount Vernon field office staff; Puget Power and Light Company: Baker River dam personnel; Skagit System Cooperative: Administration Department, spawning ground survey crews, and tagging crews; landowners who gave access to their land; and buyers and fishermen who cooperated with our efforts to recover tags.

REFERENCES CITED

- Conover, W. J. 1980. *Practical Nonparametric Statistics*. John Wiley and Sons, Inc. New York. 493 p.
- Conrad, R. H., R. A. Hayman, E. A. Beamer, and P. J. Goddard. 1993. Estimates of the number of coho salmon in the escapement to the Skagit River using a redd-count method: 1986-1990. Draft report. Northwest Indian Fisheries Commission. Olympia, WA. 112 p.
- Conrad, R. H., R. A. Hayman, E. A. Beamer, and P. J. Goddard. 1997. Coho salmon escapement to the Skagit River estimated using a mark-recapture method: 1986. Northwest Fishery Resource Bulletin, Project Report Series No. 6. Northwest Indian Fisheries Commission. Olympia, WA. 79 p.
- Conrad, R. H., R. A. Hayman, E. A. Beamer, and P. J. Goddard. 1998. Coho salmon escapement to the Skagit River estimated using a mark-recapture method: 1987. Northwest Fishery Resource Bulletin, Project Report Series No. 7. Northwest Indian Fisheries Commission. Olympia, WA. 75 p.
- Darroch, J. N. 1961. The two-sample capture-recapture census when tagging and sampling are stratified. *Biometrika* 48:241-260.
- Eames, M., T. Quinn, K. Reidinger, and M. Hino. 1981. Northern Puget Sound 1976 adult coho and chum tagging studies. Tech. Report No. 64. Washington Dept. of Fish and Wildlife. Olympia, WA. 217 p.
- Eames, M. J., T. J. Quinn, and M. Hino. 1983. 1977 Northern Puget Sound adult coho and chum tagging studies. Tech. Report No. 75. Washington Dept. of Fish and Wildlife. Olympia, WA. 239 p.
- Flint, T. 1983. Methods for estimation of Puget Sound coho salmon escapements. Unprocessed report. Washington Dept. of Fish and Wildlife. Olympia, WA. 28 p.
- Hayman, R. A. 1996. The timing and composition of the 1986-1990 Skagit coho runs. Draft report. Skagit System Cooperative. LaConner, WA.
- Hayman, R. A. 1997. Memo from R. Hayman (SSC) to Jim Seger (Pacific Fishery Management Council) dated September 25, 1997. Skagit System Cooperative. LaConner, WA.
- Johnson, R. 1986. Assessment of the Skagit River System's coho rearing potential. Unprocessed report. Washington Dept. of Fish and Wildlife. Olympia, WA.

REFERENCES CITED (continued)

- Junge, C. O. 1963. A quantitative evaluation of the bias in population estimates based on selective samples. *In* North Atlantic Fish Marking Symposium, I.C.N.A.F., Special Publication No. 4. p. 26-28.
- Mundy, P. R. 1982. Computation of migratory timing statistics for adult chinook salmon in the Yukon River, Alaska and their relevance to fisheries management. *N. Am. J. Fish. Manage.* 4:359-370.
- Noreen, E. W. 1989. *Computer Intensive Methods for Testing Hypotheses*. John Wiley and Sons, Inc. New York. 229 p.
- Pacific Fishery Management Council (PFMC). 1986. Review of 1985 Ocean Salmon Fisheries. Report funded by NOAA/NMFS Cooperative Agreement No. 86-ABH-00004. Pacific Fishery Management Council. Portland, OR.
- Pacific Fishery Management Council (PFMC). 1988. Review of 1987 Ocean Salmon Fisheries. Report funded by NOAA/NMFS Cooperative Agreement No. 88-ABH-00001. Pacific Fishery Management Council. Portland, OR.
- Pacific Fishery Management Council (PFMC). 1992. Assessment of the status of five stocks of Puget Sound chinook and coho as required under the PFMC definition of overfishing. Technical Report for the Pacific Fisheries Management Council prepared by the Puget Sound Salmon Stock Review Group. Pacific Fishery Management Council. Portland, OR. 113 p.
- Ricker, W. E. 1975. *Computation and Interpretation of Biological Statistics of Fish Populations*. Fish. Res. Board of Canada, Bull. 191. 382 p.
- Robson, D. S. and H. A. Regier. 1964. Sample size in Petersen mark-recapture experiments. *Trans. Amer. Fish. Soc.* 93:215-226.
- Seber, G. A. F. 1982. *The Estimation of Animal Abundance*. MacMillan Publishing Co. New York. 654 p.
- Washington Department of Fisheries (WDF). 1992. Washington State Sport Catch Report 1988. Washington Dept. of Fish and Wildlife. Olympia, WA. 83 p.

APPENDIX A

Summary tables of sample data for 1988.

Appendix Table A-1. Summary of coho salmon escapement samples collected at Marblemount Hatchery in 1988.

Sample Date	Sample Method	Number of Fish Examined	Number of Tags Found ^a	% with Tags (p)
19-Oct	Surplussed	857	16	1.9%
7-Nov	Pond Mortality	14	0	0.0%
10-Nov	Surplussed	25	0	0.0%
14-Nov	Pond Mortality	15	0	0.0%
15-Nov	Pond Mortality	15	1	6.7%
	Spawned	1,546	65	4.2%
	Total	1,561	66	4.2%
22-Nov	Pond Mortality	6	1	16.7%
	Spawned	1,535	62	4.0%
	Total	1,541	63	4.1%
28-Nov	Pond Mortality	40	1	2.5%
29-Nov	Pond Mortality	46	2	4.3%
	Spawned	1,242	40	3.2%
	Total	1,288	42	3.3%
30-Nov	Pond Mortality	56	2	3.6%
	Spawned	1,029	31	3.0%
	Total	1,085	33	3.0%
6-Dec	Pond Mortality	8	0	0.0%
	Spawned	773	12	1.6%
	Total	781	12	1.5%
7-Dec	Pond Mortality	164	3	1.8%
13-Dec	Pond Mortality	23	0	0.0%
	Spawned	31	0	0.0%
	Total	54	0	0.0%
20-Dec	Pond Mortality	146	4	2.7%
	Spawned	790	16	2.0%
	Total	936	20	2.1%
27-Dec	Pond Mortality	142	2	1.4%
28-Dec	Pond Mortality	12	2	16.7%
4-Jan	Pond Mortality	96	1	1.0%
	Surplussed	137	1	0.7%
	Total	233	2	0.9%
17-Jan	Pond Mortality	35	1	2.9%
Subtotals	Pond Mortality	818	20	2.4%
	Surplussed	1,019	17	1.7%
	Spawned	6,946	226	3.3%
IN-SAMPLE TOTAL		8,783	263	3.0%

^a Includes fish recovered with no tag but having the secondary mark (an opercula punch) or having an illegible tag.

Appendix Table A-2. Summary of coho salmon escapement samples collected at Baker River trap in 1988.

Sample Date	Number of Fish Examined	Number of Tags Found ^a	% with Tags (p)
9-Sep	12	0	0.0%
13-Sep	27	0	0.0%
Subtotal	39	0	0.0%
16-Sep	16	0	0.0%
19-Sep	32	0	0.0%
21-Sep	9	0	0.0%
23-Sep	22	1	4.5%
26-Sep	30	0	0.0%
28-Sep	38	0	0.0%
30-Sep	79	0	0.0%
3-Oct	112	3	2.7%
5-Oct	108	0	0.0%
7-Oct	77	1	1.3%
10-Oct	99	0	0.0%
12-Oct	105	0	0.0%
14-Oct	67	0	0.0%
17-Oct	34	1	2.9%
19-Oct	60	0	0.0%
21-Oct	45	3	6.7%
24-Oct	229	8	3.5%
26-Oct	319	7	2.2%
28-Oct	163	8	4.9%
31-Oct	124	6	4.8%
2-Nov	139	9	6.5%
4-Nov	123	8	6.5%
7-Nov	138	2	1.4%
10-Nov	92	6	6.5%
14-Nov	115	3	2.6%
16-Nov	34	3	8.8%
18-Nov	51	1	2.0%
21-Nov	60	2	3.3%
23-Nov	40	0	0.0%
28-Nov	52	1	1.9%
2-Dec	24	0	0.0%
6-Dec	9	0	0.0%
9-Dec	22	0	0.0%
13-Dec	54	1	1.9%
16-Dec	13	1	7.7%
20-Dec	7	1	14.3%
23-Dec	8	1	12.5%
30-Dec	18	0	0.0%
6-Jan	5	0	0.0%
13-Jan	3	0	0.0%
20-Jan	5	0	0.0%
27-Jan	1	0	0.0%
IN-SAMPLE TOTAL	2,781	77	2.8%

^a Includes fish recovered with no tag but having the secondary mark (an opercula punch) or having an illegible tag.

Appendix Table A-4. Summary of coho salmon catch samples collected during test fisheries in the Skagit River, 1988.

Sample Number Date	Area 2		Spudhouse		Blake's		Bay and Jetty	
	Tags Exam.	Found ^a	Fish Exam.	Tags Found ^a	Fish Exam.	Tags Found ^a	Fish Exam.	Tags Found ^a
August	20	0	139	0	43	0		
1-Sep		0.0%		0.0%		0.0%		
6-Sep			48	0	33	0		
7-Sep	25	0		0.0%	23	0		
Subtotal	45	0	187	0	99	0		
14-Sep	13	0	29	0	21	0		
15-Sep		0.0%		0.0%		0.0%		
21-Sep	9	0			58	0		
22-Sep		0.0%		0.0%		0.0%		
28-Sep	76	0	58	0	68	0		
29-Sep		0.0%		0.0%		0.0%		
5-Oct	47	0	173	0	62	0		
6-Oct		0.0%		0.0%		0.0%		
12-Oct	32	0	123	0	57	0		
13-Oct		0.0%		0.0%		0.0%		
19-Oct	124	1	325	0	266	0		
20-Oct		0.8%		0.0%		0.0%		
26-Oct	237	0	145	0			26	0
27-Oct		0.0%		0.0%		0.0%		0.0%
28-Oct					69	0		
3-Nov			19	0			53	0
4-Nov				0.0%	35	1		0.0%
IN-SAMPLE TOTAL	538	1	1,023	0	636	1	79	0
		0.2%		0.0%		0.2%		0.0%

^a Includes fish recovered with no tag but having the secondary mark (an opercula punch) or having an illegible tag.

Appendix Table A-5. Summary of coho salmon escapement samples from the Middle Skagit sub-basin collected during spawning ground surveys by Skagit System Cooperative crews and at a trap on Careys Creek, 1988.

<u>SURVEYS</u>				<u>CAREYS CREEK TRAP</u>			<u>SAMPLES COMBINED</u>		
Survey Date	Number Examined	Tags Found ^a	ρ	Number Examined	Tags Found ^a	ρ	Number Examined	Tags Found ^a	ρ
20-Oct	0	0	0.0%	15	1	6.7%	0	0	0.0%
25-Oct	0	0	0.0%				0	0	0.0%
1-Nov	0	0	0.0%				0	0	0.0%
3-Nov	0	0	0.0%				0	0	0.0%
8-Nov	0	0	0.0%				0	0	0.0%
15-Nov	2	0	0.0%				2	0	0.0%
16-Nov	3	0	0.0%				3	0	0.0%
17-Nov	1	0	0.0%				1	0	0.0%
18-Nov	3	0	0.0%				3	0	0.0%
23-Nov	7	0	0.0%				22	1	4.5%
29-Nov	4	0	0.0%				4	0	0.0%
30-Nov	19	0	0.0%				19	0	0.0%
1-Dec	11	0	0.0%				11	0	0.0%
2-Dec	12	0	0.0%				12	0	0.0%
6-Dec	2	0	0.0%				2	0	0.0%
7-Dec	38	0	0.0%				38	0	0.0%
8-Dec	64	0	0.0%				64	0	0.0%
12-Dec	53	1	1.9%				53	1	1.9%
13-Dec	6	0	0.0%				6	0	0.0%
14-Dec	21	1	4.8%				21	1	4.8%
15-Dec	38	0	0.0%				38	0	0.0%
16-Dec	17	1	5.9%				17	1	5.9%
19-Dec	21	0	0.0%				21	0	0.0%
20-Dec	15	0	0.0%				15	0	0.0%
21-Dec	2	0	0.0%				2	0	0.0%
22-Dec	20	0	0.0%				20	0	0.0%
23-Dec	79	0	0.0%				79	0	0.0%
27-Dec	45	0	0.0%				45	0	0.0%
28-Dec	56	1	1.8%				56	1	1.8%
29-Dec	25	0	0.0%				25	0	0.0%
3-Jan	1	0	0.0%				1	0	0.0%
5-Jan	84	0	0.0%				84	0	0.0%
6-Jan	25	0	0.0%				25	0	0.0%
9-Jan	0	0	0.0%				0	0	0.0%
11-Jan	0	0	0.0%				0	0	0.0%
12-Jan	40	0	0.0%				40	0	0.0%
13-Jan	17	0	0.0%				17	0	0.0%
20-Jan	43	0	0.0%				43	0	0.0%
23-Jan	13	0	0.0%				13	0	0.0%
25-Jan	0	0	0.0%				0	0	0.0%
26-Jan	4	0	0.0%				4	0	0.0%
27-Jan	39	0	0.0%				39	0	0.0%
7-Feb	0	0	0.0%				0	0	0.0%
13-Feb	3	0	0.0%				3	0	0.0%
16-Feb	0	0	0.0%				0	0	0.0%
21-Feb	0	0	0.0%				0	0	0.0%
22-Feb	0	0	0.0%				0	0	0.0%
23-Feb	0	0	0.0%				0	0	0.0%
IN-SAMPLE TOTAL	833	4	0.5%	15	1	6.7%	848	5	0.6%

^a Includes fish recovered with no tag but having the secondary mark (an opercula punch) or having an illegible tag.

Appendix Table A-6. Summary of coho salmon escapement samples from the Middle Skagit sub-basin collected at a trap on Etach Slough operated by Skagit System Cooperative, 1988.

<u>RACK RECOVERIES</u>				<u>TRAP RECOVERIES</u>			<u>SAMPLES COMBINED</u>		
Date	Number Examined	Tags Found ^a	ρ	Number Examined	Tags Found ^a	ρ	Number Examined	Tags Found ^a	ρ
1-Nov				2	0	0.0%	2	0	0.0%
3-Nov				1	0	0.0%	1	0	0.0%
4-Nov				1	0	0.0%	1	0	0.0%
5-Nov				2	0	0.0%	2	0	0.0%
7-Nov				9	1	11.1%	9	1	11.1%
8-Nov				2	0	0.0%	2	0	0.0%
9-Nov				2	0	0.0%	2	0	0.0%
10-Nov				3	1	33.3%	3	1	33.3%
11-Nov				7	1	14.3%	7	1	14.3%
12-Nov				6	1	16.7%	6	1	16.7%
13-Nov				1	0	0.0%	1	0	0.0%
14-Nov	0	0	0.0%	2	0	0.0%	2	0	0.0%
16-Nov				1	0	0.0%	1	0	0.0%
17-Nov				1	0	0.0%	1	0	0.0%
20-Nov	1	1	100.0%	21	1	4.8%	22	2	9.1%
21-Nov				16	0	0.0%	16	0	0.0%
22-Nov				9	0	0.0%	9	0	0.0%
23-Nov	1	1	100.0%	1	0	0.0%	2	1	50.0%
24-Nov	3	0	0.0%	5	0	0.0%	8	0	0.0%
25-Nov				2	0	0.0%	2	0	0.0%
26-Nov				1	0	0.0%	1	0	0.0%
27-Nov	2	0	0.0%				2	0	0.0%
28-Nov				6	0	0.0%	6	0	0.0%
29-Nov	2	0	0.0%	6	0	0.0%	8	0	0.0%
30-Nov	2	0	0.0%	4	0	0.0%	6	0	0.0%
Subtotal	11	2	18.2%	111	5	4.5%	122	7	5.7%
1-Dec	1	0	0.0%	5	0	0.0%	6	0	0.0%
2-Dec	2	0	0.0%	2	0	0.0%	4	0	0.0%
3-Dec				8	0	0.0%	8	0	0.0%
4-Dec	0	0	0.0%	1	0	0.0%	1	0	0.0%
5-Dec	0	0	0.0%	2	0	0.0%	2	0	0.0%
6-Dec	3	0	0.0%	6	1	16.7%	9	1	11.1%
7-Dec	2	0	0.0%	5	0	0.0%	7	0	0.0%
8-Dec	0	0	0.0%	2	0	0.0%	2	0	0.0%
9-Dec	0	0	0.0%	3	0	0.0%	3	0	0.0%
10-Dec	1	0	0.0%	1	0	0.0%	2	0	0.0%
11-Dec	2	0	0.0%	1	0	0.0%	3	0	0.0%
12-Dec				6	0	0.0%	6	0	0.0%
15-Dec				4	0	0.0%	4	0	0.0%
16-Dec				2	0	0.0%	2	0	0.0%
17-Dec				2	0	0.0%	2	0	0.0%
18-Dec				4	0	0.0%	4	0	0.0%
19-Dec	1	0	0.0%	2	0	0.0%	3	0	0.0%
20-Dec				1	0	0.0%	1	0	0.0%
21-Dec	0	0	0.0%	3	1	33.3%	3	1	33.3%
22-Dec	0	0	0.0%				0	0	0.0%
23-Dec				1	0	0.0%	1	0	0.0%
24-Dec				3	0	0.0%	3	0	0.0%

- continued -

Appendix Table A-6. Summary of coho salmon escapement samples from the Middle Skagit sub-basin collected at a trap on Etach Slough operated by Skagit System Cooperative, 1988 (continued).

<u>RACK RECOVERIES</u>				<u>TRAP RECOVERIES</u>			<u>SAMPLES COMBINED</u>		
Survey	Number	Tags		Number	Tags		Number	Tags	
Date	Examined	Found ^a	ρ	Examined	Found ^a	ρ	Examined	Found ^a	ρ
25-Dec	1	0	0.0%				1	0	0.0%
26-Dec	0	0	0.0%				0	0	0.0%
27-Dec				2	0	0.0%	2	0	0.0%
28-Dec				1	0	0.0%	1	0	0.0%
29-Dec	1	0	0.0%	1	0	0.0%	2	0	0.0%
30-Dec	0	0	0.0%	20	0	0.0%	20	0	0.0%
31-Dec	0	0	0.0%	6	0	0.0%	6	0	0.0%
1-Jan				7	0	0.0%	7	0	0.0%
2-Jan	1	0	0.0%	6	0	0.0%	7	0	0.0%
3-Jan	0	0	0.0%	4	0	0.0%	4	0	0.0%
4-Jan	0	0	0.0%	56	0	0.0%	56	0	0.0%
5-Jan				6	0	0.0%	6	0	0.0%
6-Jan				1	0	0.0%	1	0	0.0%
7-Jan				4	0	0.0%	4	0	0.0%
9-Jan	0	0	0.0%	3	0	0.0%	3	0	0.0%
10-Jan	0	0	0.0%	3	0	0.0%	3	0	0.0%
11-Jan	0	0	0.0%	4	0	0.0%	4	0	0.0%
12-Jan				6	0	0.0%	6	0	0.0%
13-Jan	0	0	0.0%	2	0	0.0%	2	0	0.0%
14-Jan				2	0	0.0%	2	0	0.0%
15-Jan	1	0	0.0%	3	1	33.3%	4	1	25.0%
16-Jan				52	0	0.0%	52	0	0.0%
17-Jan	0	0	0.0%	22	0	0.0%	22	0	0.0%
18-Jan				6	0	0.0%	6	0	0.0%
19-Jan	1	0	0.0%	1	0	0.0%	2	0	0.0%
20-Jan	0	0	0.0%	1	0	0.0%	1	0	0.0%
21-Jan	0	0	0.0%	1	0	0.0%	1	0	0.0%
22-Jan	0	0	0.0%				0	0	0.0%
23-Jan	0	0	0.0%				0	0	0.0%
24-Jan	0	0	0.0%	1	0	0.0%	1	0	0.0%
25-Jan	0	0	0.0%	1	0	0.0%	1	0	0.0%
26-Jan	0	0	0.0%				0	0	0.0%
29-Jan	0	0	0.0%				0	0	0.0%
31-Jan				1	0	0.0%	1	0	0.0%
3-Feb	0	0	0.0%				0	0	0.0%
4-Feb				1	0	0.0%	1	0	0.0%
11-Feb	0	0	0.0%				0	0	0.0%
Subtotal	17	0	0.0%	288	3	1.0%	305	3	1.0%
IN-SAMPLE TOTAL	28	2	7.1%	399	8	2.0%	427	10	2.3%

^a Includes fish recovered with no tag but having the secondary mark (an opercula punch) or having an illegible tag.

Appendix Table A-7. Summary of coho salmon escapement samples from the Upper Skagit sub-basin collected during spawning ground surveys by Skagit System Cooperative crews, 1988.

Survey Date	Number of Fish Examined	Number of Tags Found ^a	% with Tags (p)
27-Oct	0	0	0.0%
31-Oct	0	0	0.0%
18-Nov	2	0	0.0%
21-Nov	4	0	0.0%
28-Nov	2	0	0.0%
29-Nov	15	0	0.0%
1-Dec	2	0	0.0%
2-Dec	5	1	20.0%
3-Dec	2	0	0.0%
5-Dec	3	1	33.3%
6-Dec	29	1	3.4%
7-Dec	1	0	0.0%
8-Dec	3	0	0.0%
9-Dec	20	1	5.0%
12-Dec	7	0	0.0%
13-Dec	17	0	0.0%
15-Dec	4	0	0.0%
16-Dec	5	0	0.0%
19-Dec	1	0	0.0%
20-Dec	21	0	0.0%
21-Dec	6	0	0.0%
22-Dec	11	0	0.0%
23-Dec	5	1	20.0%
27-Dec	14	2	14.3%
28-Dec	2	0	0.0%
29-Dec	20	0	0.0%
30-Dec	8	0	0.0%
5-Jan	40	1	2.5%
6-Jan	2	0	0.0%
12-Jan	6	0	0.0%
13-Jan	1	0	0.0%
16-Jan	63	1	1.6%
18-Jan	0	0	0.0%
19-Jan	11	1	9.1%
20-Jan	4	0	0.0%
25-Jan	9	0	0.0%
26-Jan	10	0	0.0%
27-Jan	25	0	0.0%
30-Jan	7	0	0.0%
6-Feb	21	0	0.0%
7-Feb	5	0	0.0%
9-Feb	7	0	0.0%
10-Feb	15	0	0.0%
14-Feb	2	0	0.0%
15-Feb	5	0	0.0%
17-Feb	19	1	5.3%
24-Feb	12	0	0.0%
1-Mar	4	0	0.0%
29-Mar	2	0	0.0%
IN-SAMPLE TOTAL	479	11	2.3%

^a Includes fish recovered with no tag but having the secondary mark (an opercula punch) or having an illegible tag.

Appendix Table A-8. Summary of coho salmon escapement samples from the Lower Sauk sub-basin collected during spawning ground surveys by Skagit System Cooperative crews, 1988.

Survey Date	Number of Fish Examined	Number of Tags Found ^a	% with Tags (p)
3-Nov	0	0	0.0%
14-Nov	0	0	0.0%
16-Nov	1	0	0.0%
28-Nov	7	0	0.0%
1-Dec	3	0	0.0%
8-Dec	10	1	10.0%
12-Dec	6	1	16.7%
15-Dec	13	0	0.0%
19-Dec	30	1	3.3%
22-Dec	4	0	0.0%
27-Dec	2	0	0.0%
29-Dec	39	1	2.6%
30-Dec	0	0	0.0%
6-Jan	12	1	8.3%
7-Jan	35	2	5.7%
13-Jan	5	1	20.0%
16-Jan	77	2	2.6%
19-Jan	1	0	0.0%
20-Jan	7	0	0.0%
26-Jan	23	0	0.0%
27-Jan	156	3	1.9%
30-Jan	4	0	0.0%
6-Feb	70	0	0.0%
7-Feb	70	1	1.4%
13-Feb	26	0	0.0%
16-Feb	22	0	0.0%
21-Feb	26	0	0.0%
27-Feb	5	0	0.0%
IN-SAMPLE TOTAL	654	14	2.1%

^a Includes fish recovered with no tag but having the secondary mark (an opercula punch) or having an illegible tag.

Appendix Table A-9. Summary of coho salmon escapement samples from the Middle Sauk sub-basin collected during spawning ground surveys by Skagit System Cooperative crews, 1988.

Survey Date	Number of Fish Examined	Number of Tags Found ^a	% with Tags (p)
22-Oct	0	0	0.0%
27-Oct	0	0	0.0%
3-Nov	0	0	0.0%
14-Nov	2	0	0.0%
18-Nov	0	0	0.0%
28-Nov	0	0	0.0%
1-Dec	5	0	0.0%
2-Dec	5	0	0.0%
9-Dec	12	0	0.0%
16-Dec	12	0	0.0%
19-Dec	63	5	7.9%
22-Dec	34	0	0.0%
29-Dec	63	0	0.0%
4-Jan	29	0	0.0%
6-Jan	35	0	0.0%
11-Jan	15	1	6.7%
13-Jan	60	0	0.0%
19-Jan	19	0	0.0%
20-Jan	61	1	1.6%
25-Jan	58	1	1.7%
27-Jan	113	1	0.9%
6-Feb	21	0	0.0%
7-Feb	10	0	0.0%
10-Feb	7	0	0.0%
13-Feb	36	1	2.8%
23-Feb	23	0	0.0%
IN-SAMPLE TOTAL	683	10	1.5%

^a Includes fish recovered with no tag but having the secondary mark (an opercula punch) or having an illegible tag.

Appendix Table A-10. Summary of coho salmon escapement samples from the Upper Sauk sub-basin collected during spawning ground surveys by Skagit System Cooperative crews, 1988.

Survey Date	Number of Fish Examined	Number of Tags Found ^a	% with Tags (p)
2-Nov	3	1	33.3%
16-Nov	2	0	0.0%
1-Dec	1	0	0.0%
8-Dec	5	0	0.0%
16-Dec	5	0	0.0%
22-Dec	5	0	0.0%
29-Dec	9	1	11.1%
4-Jan	3	0	0.0%
18-Jan	1	0	0.0%
24-Jan	9	1	11.1%
10-Feb	5	1	20.0%
15-Feb	2	0	0.0%
IN-SAMPLE TOTAL	50	4	8.0%

^a Includes fish recovered with no tag but having the secondary mark (an opercula punch) or having an illegible tag.

Appendix Table A-11. Summary of coho salmon escapement samples from the Suiattle sub-basin collected during spawning ground surveys by Skagit System Cooperative crews and at a trap on Suiattle Slough, 1988.

<u>SURVEYS</u>				<u>SUIATTLE TRAP</u>			<u>SAMPLES COMBINED</u>		
Survey Date	Number Examined	Tags Found ^a	ρ	Number Examined	Tags Found ^a	ρ	Number Examined	Tags Found ^a	ρ
14-Nov	1	0	0.0%				1	0	0.0%
21-Nov	2	0	0.0%				2	0	0.0%
26-Nov				2	0	0.0%	2	0	0.0%
28-Nov	4	0	0.0%				4	0	0.0%
5-Dec	10	0	0.0%	1	0	0.0%	11	0	0.0%
7-Dec				3	0	0.0%	3	0	0.0%
9-Dec				1	0	0.0%	1	0	0.0%
12-Dec				2	0	0.0%	2	0	0.0%
13-Dec				3	0	0.0%	3	0	0.0%
14-Dec	18	1	5.6%	1	0	0.0%	19	1	5.3%
16-Dec				1	0	0.0%	1	0	0.0%
21-Dec	46	1	2.2%	2	0	0.0%	48	1	2.1%
27-Dec				3	0	0.0%	3	0	0.0%
28-Dec	101	2	2.0%				101	2	2.0%
30-Dec				22	1	4.5%	22	1	4.5%
1-Jan				6	1	16.7%	6	1	16.7%
3-Jan				3	0	0.0%	3	0	0.0%
4-Jan				19	0	0.0%	19	0	0.0%
5-Jan				13	0	0.0%	13	0	0.0%
6-Jan	87	3	3.4%	1	0	0.0%	88	3	3.4%
9-Jan				4	0	0.0%	4	0	0.0%
10-Jan				4	0	0.0%	4	0	0.0%
11-Jan				2	0	0.0%	2	0	0.0%
13-Jan				6	0	0.0%	6	0	0.0%
15-Jan				6	0	0.0%	6	0	0.0%
17-Jan	86	1	1.2%	20	0	0.0%	106	1	0.9%
18-Jan				13	0	0.0%	13	0	0.0%
19-Jan	8	1	12.5%	2	0	0.0%	10	1	10.0%
20-Jan				2	0	0.0%	2	0	0.0%
23-Jan	51	2	3.9%	5	0	0.0%	56	2	3.6%
24-Jan				3	0	0.0%	3	0	0.0%
25-Jan				3	0	0.0%	3	0	0.0%
27-Jan				9	0	0.0%	9	0	0.0%
30-Jan	72	1	1.4%	6	0	0.0%	78	1	1.3%
31-Jan				16	0	0.0%	16	0	0.0%
6-Feb				1	0	0.0%	1	0	0.0%
9-Feb	3	0	0.0%				3	0	0.0%
14-Feb	5	0	0.0%				5	0	0.0%
IN-SAMPLE TOTAL	494	12	2.4%	185	2	1.1%	679	14	2.1%

^a Includes fish recovered with no tag but having the secondary mark (an opercula punch) or having an illegible tag.

Appendix Table A-12. Summary of coho salmon escapement samples from the Nookachamps sub-basin collected during spawning ground surveys by Skagit System Cooperative crews and at traps on the East Fork and West Fork of Nookachamps Creek, 1988.

Survey Date	SSCSURVEYS			EAST FORK TRAP			WEST FORK TRAP			SAMPLES COMBINED		
	Number Exam.	Tags Found ^a	% w/Tags (ρ)	Fish Exam.	Tags Found ^a	% w/Tags (ρ)	Fish Exam.	Tags Found ^a	% w/Tags (ρ)	Fish Exam.	Tags Found ^a	% w/Tags (ρ)
23-Oct				2	0	0.0%	1	0	0.0%	2	0	0.0%
25-Oct										1	0	0.0%
28-Oct				1	0	0.0%				1	0	0.0%
1-Nov	0	0	0.0%	1	0	0.0%				1	0	0.0%
2-Nov				16	0	0.0%				16	0	0.0%
3-Nov				3	0	0.0%				3	0	0.0%
4-Nov							1	0	0.0%	1	0	0.0%
5-Nov							2	0	0.0%	2	0	0.0%
6-Nov				6	0	0.0%				6	0	0.0%
7-Nov							1	0	0.0%	1	0	0.0%
8-Nov	0	0	0.0%	1	0	0.0%	1	0	0.0%	2	0	0.0%
17-Nov	12	0	0.0%	1	0	0.0%	2	0	0.0%	15	0	0.0%
18-Nov				1	0	0.0%				1	0	0.0%
19-Nov				1	0	0.0%				1	0	0.0%
21-Nov				1	0	0.0%	3	0	0.0%	4	0	0.0%
29-Nov	16	0	0.0%							16	0	0.0%
30-Nov	9	0	0.0%							9	0	0.0%
1-Dec				1	0	0.0%				1	0	0.0%
2-Dec	24	0	0.0%	1	0	0.0%				25	0	0.0%
6-Dec	4	0	0.0%							4	0	0.0%
7-Dec	34	0	0.0%							34	0	0.0%
8-Dec	67	1	1.5%	1	0	0.0%				68	1	1.5%
9-Dec				2	0	0.0%				2	0	0.0%
10-Dec				1	0	0.0%				1	0	0.0%
12-Dec	7	0	0.0%							7	0	0.0%
13-Dec				7	0	0.0%				7	0	0.0%
14-Dec				1	0	0.0%				1	0	0.0%
15-Dec				1	0	0.0%				1	0	0.0%

- continued -

Appendix Table A-12. Summary of coho salmon escapement samples from the Nookachamps sub-basin collected during spawning ground surveys by Skagit System Cooperative crews and at traps on the East Fork and West Fork of Nookachamps Creek, 1988 (continued).

Survey Date	SSCSURVEYS			EAST FORK TRAP			WEST FORK TRAP			SAMPLES COMBINED		
	Number	Tags	% w/Tags (ρ)	Fish Exam.	Tags Found ^a	% w/Tags (ρ)	Fish Exam.	Tags Found ^a	% w/Tags (ρ)	Fish Exam.	Tags Found ^a	% w/Tags (ρ)
16-Dec	43	0	0.0%							43	0	0.0%
20-Dec	45	0	0.0%							45	0	0.0%
23-Dec	16	0	0.0%							16	0	0.0%
27-Dec	8	0	0.0%							8	0	0.0%
28-Dec	24	0	0.0%							24	0	0.0%
29-Dec	24	0	0.0%							24	0	0.0%
30-Dec				2	0	0.0%				2	0	0.0%
2-Jan				1	0	0.0%				1	0	0.0%
5-Jan	12	0	0.0%							12	0	0.0%
12-Jan	7	0	0.0%							7	0	0.0%
27-Jan	4	0	0.0%							4	0	0.0%
IN-SAMPLE TOTAL	356	1	0.3%	52	0	0.0%	11	0	0.0%	419	1	0.2%

^a Includes fish recovered with no tag but having the secondary mark (an opercula punch) or having an illegible tag.

Appendix Table A-13. Summary of coho salmon escapement samples from the Carpenter sub-basin collected during spawning ground surveys by Skagit System Cooperative crews and at a trap on Carpenter Creek Slough, 1988.

SURVEYS				SLOUGH TRAP			SAMPLES COMBINED		
Survey Date	Number Examined	Tags Found ^a	ρ	Number Examined	Tags Found ^a	ρ	Number Examined	Tags Found ^a	ρ
4-Nov				1	0	0.0%	1	0	0.0%
6-Nov				135	0	0.0%	135	0	0.0%
7-Nov				115	0	0.0%	115	0	0.0%
8-Nov	2	0	0.0%	30	0	0.0%	32	0	0.0%
9-Nov				1	0	0.0%	1	0	0.0%
11-Nov				1	0	0.0%	1	0	0.0%
13-Nov				7	0	0.0%	7	0	0.0%
14-Nov				1	0	0.0%	1	0	0.0%
16-Nov				1	0	0.0%	1	0	0.0%
21-Nov				5	0	0.0%	5	0	0.0%
22-Nov				143	0	0.0%	143	0	0.0%
23-Nov				80	0	0.0%	80	0	0.0%
24-Nov				17	0	0.0%	17	0	0.0%
25-Nov				4	0	0.0%	4	0	0.0%
26-Nov				3	0	0.0%	3	0	0.0%
27-Nov				0	0	0.0%	0	0	0.0%
28-Nov				2	0	0.0%	2	0	0.0%
29-Nov				24	0	0.0%	24	0	0.0%
30-Nov				5	0	0.0%	5	0	0.0%
1-Dec	0	0	0.0%	1	0	0.0%	1	0	0.0%
5-Dec	13	0	0.0%				13	0	0.0%
6-Dec				1	0	0.0%	1	0	0.0%
7-Dec	3	0	0.0%	8	0	0.0%	11	0	0.0%
8-Dec				3	0	0.0%	3	0	0.0%
9-Dec	77	0	0.0%	4	0	0.0%	81	0	0.0%
10-Dec				2	0	0.0%	2	0	0.0%
11-Dec				1	0	0.0%	1	0	0.0%
12-Dec	0	0	0.0%	33	0	0.0%	33	0	0.0%
13-Dec	2	0	0.0%	13	0	0.0%	15	0	0.0%
14-Dec				10	0	0.0%	10	0	0.0%
15-Dec	19	0	0.0%	4	0	0.0%	23	0	0.0%
16-Dec				1	0	0.0%	1	0	0.0%
17-Dec	2	0	0.0%				2	0	0.0%
19-Dec				2	0	0.0%	2	0	0.0%
20-Dec				1	0	0.0%	1	0	0.0%
21-Dec	2	0	0.0%				2	0	0.0%
22-Dec	23	0	0.0%				23	0	0.0%
23-Dec	0	0	0.0%				0	0	0.0%
26-Dec				0	0	0.0%	0	0	0.0%
30-Dec				2	0	0.0%	2	0	0.0%
31-Dec				1	0	0.0%	1	0	0.0%
1-Jan				1	0	0.0%	1	0	0.0%
2-Jan				3	0	0.0%	3	0	0.0%
3-Jan	2	0	0.0%	0	0	0.0%	2	0	0.0%
12-Jan	0	0	0.0%				0	0	0.0%
13-Jan	27	0	0.0%				27	0	0.0%
IN-SAMPLE TOTAL	172	0	0.0%	666	0	0.0%	838	0	0.0%

^a Includes fish recovered with no tag but having the secondary mark (an opercula punch) or having an illegible tag.

Appendix Table A-14. Summary of coho salmon escapement samples from the Cascade sub-basin collected during spawning ground surveys by Skagit System Cooperative crews, 1988.

Survey Date	Number of Fish Examined	Number of Tags Found ^a	% with Tags (p)
7-Nov	1	0	0.0%
14-Nov	1	0	0.0%
17-Nov	14	0	0.0%
19-Nov	3	0	0.0%
28-Nov	6	0	0.0%
30-Nov	9	0	0.0%
5-Dec	11	0	0.0%
6-Dec	21	0	0.0%
12-Dec	11	0	0.0%
16-Dec	5	0	0.0%
19-Dec	19	1	5.3%
20-Dec	1	1	100.0%
27-Dec	5	0	0.0%
28-Dec	22	0	0.0%
4-Jan	16	0	0.0%
9-Jan	23	3	13.0%
10-Jan	1	1	100.0%
17-Jan	1	0	0.0%
18-Jan	5	0	0.0%
24-Jan	0	0	0.0%
25-Jan	4	1	25.0%
26-Jan	2	0	0.0%
30-Jan	4	0	0.0%
7-Feb	2	0	0.0%
14-Feb	4	0	0.0%
22-Feb	0	0	0.0%
23-Feb	0	0	0.0%
IN-SAMPLE TOTAL	191	7	3.7%

^a Includes fish recovered with no tag but having the secondary mark (an opercula punch) or having an illegible tag.

Appendix Table A-15. CPUE (catch per beach seine set) of coho salmon bound for major recovery areas in the Skagit River, 1988. CPUE for recovery areas estimated using in-sample tag recoveries.

Recoveries by release strata.						
Tag Release Period	Number of Sets	Coho Catch	Catch/ Set	MM Hatchery	Baker R. Trap	Middle Skagit
1. 01-Sep to 10-Sep	0	0		0	0	0
2. 11-Sep to 20-Sep	58	264	4.6	15	5	0
3. 21-Sep to 30-Sep	33	133	4.0	9	5	0
4. 01-Oct to 10-Oct	50	657	13.1	76	19	3
5. 11-Oct to 20-Oct	42	607	14.5	80	23	5
6. 21-Oct to 30-Oct	43	407	9.5	42	14	2
7. 31-Oct to 09-Nov	34	90	2.6	6	1	0
8. 10-Nov to 19-Nov	23	82	3.6	3	0	0
9. 20-Nov to 29-Nov	0	0		0	0	0
10. 30-Nov to 09-Dec	0	0		0	0	0
Totals	283	2,240	7.9	231	67	10
CPUE of fish bound for indicated recovery areas.						
				Release Period	MM Hatchery	Baker R. Trap
						Middle Skagit
				1	0.00	0.00
				2	0.26	0.09
				3	0.27	0.15
				4	1.52	0.38
				5	1.90	0.55
				6	0.98	0.33
				7	0.18	0.03
				8	0.13	0.00
				9	0.00	0.00
				10	0.00	0.00
				Totals	5.24	1.52
CPUE standardized as a percentage of total for area.						
				Release Period	MM Hatchery	Baker R. Trap
						Middle Skagit
				1	0.0%	0.0%
				2	4.9%	5.7%
				3	5.2%	10.0%
				4	29.0%	25.0%
				5	36.4%	36.0%
				6	18.6%	21.4%
				7	3.4%	1.9%
				8	2.5%	0.0%
				9	0.0%	0.0%
				10	0.0%	0.0%
				Totals	100.0%	100.0%

Appendix Table A-16. Summary of the number of tag releases and number of in-sample tag recoveries by length for male and female coho salmon tagged in the lower Skagit River, 1988.

Length in cm	MALES			FEMALES		
	Number Released	Number Recovered	Percent Recovered	Number Released	Number Recovered	Percent Recovered
≤ 35	10	0	0.0%	0	0	0.0%
36	8	1	12.5%	0	0	0.0%
37	5	0	0.0%	0	0	0.0%
38	6	1	16.7%	1	0	0.0%
39	8	0	0.0%	1	0	0.0%
40	14	0	0.0%	0	0	0.0%
41	22	2	9.1%	0	0	0.0%
42	26	0	0.0%	0	0	0.0%
43	36	2	5.6%	0	0	0.0%
44	31	5	16.1%	0	0	0.0%
45	35	4	11.4%	3	2	66.7%
46	63	6	9.5%	1	1	100.0%
47	51	7	13.7%	2	1	50.0%
48	58	12	20.7%	10	0	0.0%
49	61	2	3.3%	13	2	15.4%
Subtotal	434	42	9.7%			
50	53	10	18.9%	16	5	31.3%
51	70	12	17.1%	18	2	11.1%
52	56	13	23.2%	35	8	22.9%
53	75	20	26.7%	26	8	30.8%
54	62	21	33.9%	41	5	12.2%
55	54	18	33.3%	45	12	26.7%
56	37	14	37.8%	63	15	23.8%
57	47	11	23.4%	59	13	22.0%
58	32	9	28.1%	70	23	32.9%
				404	97	24.0%
59	31	9	29.0%	47	6	12.8%
60	36	10	27.8%	48	9	18.8%
61	32	11	34.4%	42	3	7.1%
62	21	5	23.8%	53	13	24.5%
63	18	6	33.3%	32	6	18.8%
64	23	4	17.4%	36	5	13.9%
65	21	5	23.8%	27	0	0.0%
66	22	7	31.8%	20	2	10.0%
67	15	3	20.0%	18	1	5.6%
68	16	3	18.8%	15	2	13.3%
69	17	5	29.4%	9	2	22.2%
70	10	1	10.0%	3	0	0.0%
71	9	2	22.2%	2	0	0.0%
72	6	3	50.0%	1	1	100.0%
73	4	0	0.0%	0	0	0.0%
74	5	0	0.0%	0	0	0.0%
75	2	0	0.0%	0	0	0.0%
76	1	0	0.0%	0	0	0.0%
77	1	0	0.0%	0	0	0.0%
78	1	0	0.0%	0	0	0.0%
79	1	0	0.0%	0	0	0.0%
80	0	0	0.0%	0	0	0.0%
>80	1	0	0.0%	0	0	0.0%
Subtotal	779	202	25.9%	353	50	14.2%
TOTAL	1,213	244	20.1%	757	147	19.4%

Appendix Table A-17. Daily summary of the numbers of coho salmon tagged in the lower Skagit River and recovered during in-sample surveys, by sex, release condition, and maturity classification, 1988.

Date	SEX				CONDITION						MATURITY					
	<u>Male</u>		<u>Female</u>		<u>X-</u>		<u>X</u>		<u>X+</u>		<u>Bright</u>		<u>Blush</u>		<u>Dark</u>	
	Rel.	Rec.	Rel.	Rec.	Rel.	Rec.	Rel.	Rec.	Rel.	Rec.	Rel.	Rec.	Rel.	Rec.	Rel.	Rec.
12-Sep	23	1	11	1	2	0	0	0	32	2	34	2	0	0	0	0
13-Sep	41	6	19	2	4	0	56	8	0	0	60	8	0	0	0	0
15-Sep	10	0	12	0	1	0	21	0	0	0	22	0	0	0	0	0
16-Sep	4	1	8	3	3	1	9	3	0	0	12	4	0	0	0	0
19-Sep	25	5	20	6	1	0	44	11	0	0	45	11	0	0	0	0
20-Sep	28	5	16	6	2	0	42	11	0	0	44	11	0	0	0	0
23-Sep	40	9	24	7	1	0	63	16	0	0	60	15	4	1	0	0
27-Sep	11	4	7	5	3	1	15	8	0	0	18	9	0	0	0	0
28-Sep	25	8	10	5	3	1	32	12	0	0	34	12	1	1	0	0
3-Oct	68	15	38	3	5	0	101	18	0	0	96	18	9	0	0	0
4-Oct	33	5	28	8	0	0	61	13	0	0	61	13	0	0	0	0
6-Oct	49	11	38	7	3	0	83	17	1	1	85	18	2	0	0	0
7-Oct	104	25	60	9	1	0	163	34	0	0	160	32	3	1	1	1
10-Oct	113	17	40	5	2	0	151	22	0	0	149	22	4	0	0	0
11-Oct	34	7	18	5	2	0	50	12	0	0	44	9	8	3	0	0
13-Oct	48	13	41	10	0	0	89	23	0	0	83	21	6	2	0	0
14-Oct	185	42	110	23	7	2	288	63	0	0	250	54	45	11	0	0
20-Oct	64	16	34	6	0	0	98	22	0	0	68	15	29	7	1	0
21-Oct	115	20	61	10	5	1	171	29	0	0	125	17	50	13	1	0
25-Oct	45	10	35	9	0	0	80	19	0	0	66	13	13	5	1	1
26-Oct	35	3	31	3	0	0	66	6	0	0	61	6	5	0	0	0
28-Oct	43	9	23	3	1	0	65	12	0	0	59	9	7	3	0	0
1-Nov	16	3	21	5	1	0	36	8	0	0	30	5	7	3	0	0
2-Nov	18	3	14	1	0	0	32	4	0	0	23	2	4	1	5	1
9-Nov	9	3	9	1	1	1	17	3	0	0	8	1	5	1	5	2
10-Nov	2	0	2	0	0	0	4	0	0	0	2	0	2	0	0	0
15-Nov	25	3	27	4	1	0	51	7	0	0	46	6	6	1	0	0
Total	1,213	244	757	147	49	7	1,888	381	33	3	1,745	333	210	53	14	5
% Recovered		20.1		19.4			14.3		20.2			19.1		25.2		35.7

Appendix Table A-18. Summary of the estimated number of tags from areas downstream of the tagging area in the lower Skagit River, 1988.

A. Downstream commercial fishery and test fishery catches.

Area	Catch ^a Before Tagging	Catch After Tagging	Number of Fish Examined	Number of Tags Found	Estimated Total Tags Present
8E	0	1,946	532	0	
8	88	1,230	300	0	
78C	794	3,657	696	0	
Test Fishery ^b	331	2,289	2,276	2	
78D-1, 78D-2	1	2,243 ^c	1,264	1	
Total	1,214	11,365	5,068	3	6.7

^a Catches prior to tagging not included in tag recovery expansions.

^b Test fisheries at Area 2, Blakes, Bay, and Jetty.

^c Estimated catch below the tagging area by the commercial fishery after the onset of tagging.

B. Downstream spawning areas (redd data from Conrad et al. [1993]).

Area	Estimated Number of Redds	Estimated Number of Fish/Redd	Estimated Total Escapement	Number of Fish Examined	Number of Tags Found	Estimated Total Tags Present
Carpenter	1,541	1.2	1,849	838	0	
Nookachamps	1,171	1.2	1,405	419	1	
Total	2,712		3,254	1,257	1	2.6

APPENDIX B

Details of abundance estimates generated for 1988.

APPENDIX B

RECOVERY LOCATION: Marblemount Hatchery

ESTIMATION METHOD: Petersen

95% CONFIDENCE INTERVAL: Normal Approximation

TAG RELEASE AND RECOVERY SUMMARY:

Number of Tags Released = 1,970

Number of Fish Examined for Tags = 8,783

Number of Tagged or Marked Fish Recovered = 263

RECOVERY LOCATION: Baker River Trap

ESTIMATION METHOD: Petersen

95% CONFIDENCE INTERVAL: Normal Approximation

TAG RELEASE AND RECOVERY SUMMARY:

Number of Tags Released = 1,970

Number of Fish Examined for Tags = 2,781

Number of Tagged or Marked Fish Recovered = 77

RECOVERY LOCATION: Marblemount-Baker Pooled

ESTIMATION METHOD: Petersen

95% CONFIDENCE INTERVAL: Normal Approximation

TAG RELEASE AND RECOVERY SUMMARY:

Number of Tags Released = 1,970

Number of Fish Examined for Tags = 11,564

Number of Tagged or Marked Fish Recovered = 340

APPENDIX B

RECOVERY LOCATION: Lower Sauk Sub-basin

ESTIMATION METHOD: Petersen

95% CONFIDENCE INTERVAL: Poisson Approximation

TAG RELEASE AND RECOVERY SUMMARY:

Number of Tags Released = 1,970

Number of Fish Examined for Tags = 654

Number of Tagged or Marked Fish Recovered = 14

RECOVERY LOCATION: Suiattle Sub-basin

ESTIMATION METHOD: Petersen

95% CONFIDENCE INTERVAL: Poisson Approximation

TAG RELEASE AND RECOVERY SUMMARY:

Number of Tags Released = 1,970

Number of Fish Examined for Tags = 679

Number of Tagged or Marked Fish Recovered = 14

RECOVERY LOCATION: Upper Skagit Sub-basin

ESTIMATION METHOD: Petersen

95% CONFIDENCE INTERVAL: Poisson Approximation

TAG RELEASE AND RECOVERY SUMMARY:

Number of Tags Released = 1,970

Number of Fish Examined for Tags = 479

Number of Tagged or Marked Fish Recovered = 11

APPENDIX B

RECOVERY LOCATION: Middle Sauk Sub-basin

ESTIMATION METHOD: Petersen

95% CONFIDENCE INTERVAL: Poisson Approximation

TAG RELEASE AND RECOVERY SUMMARY:

Number of Tags Released = 1,970

Number of Fish Examined for Tags = 683

Number of Tagged or Marked Fish Recovered = 10

RECOVERY LOCATION: Cascade Sub-basin

ESTIMATION METHOD: Petersen

95% CONFIDENCE INTERVAL: Poisson Approximation

TAG RELEASE AND RECOVERY SUMMARY:

Number of Tags Released = 1,970

Number of Fish Examined for Tags = 191

Number of Tagged or Marked Fish Recovered = 7

RECOVERY LOCATION: Upper Skagit-Lower Sauk-Middle Sauk-Suiattle-
Cascade Pooled

ESTIMATION METHOD: Petersen

95% CONFIDENCE INTERVAL: Normal Approximation

TAG RELEASE AND RECOVERY SUMMARY:

Number of Tags Released = 1,970

Number of Fish Examined for Tags = 2,686

Number of Tagged or Marked Fish Recovered = 56

APPENDIX B

RECOVERY LOCATION: Marblemount-Baker-Upper Skagit-Lower Sauk-Middle Sauk-Suiattle-Cascade Pooled

ESTIMATION METHOD: Petersen

95% CONFIDENCE INTERVAL: Normal Approximation

TAG RELEASE AND RECOVERY SUMMARY:

Number of Tags Released = 1,970

Number of Fish Examined for Tags = 14,250

Number of Tagged or Marked Fish Recovered = 396
